

Guidelines for the Prevention of Intravascular Catheter-Related Infections

Prepared by

Naomi P. O'Grady, M.D.¹
Mary Alexander, R.N.²
Lillian A. Burns, M.T., M.P.H., C.I.C.³
E. Patchen Dellinger, M.D.⁴
Jeffery Garland, M.D.⁵
Stephen O. Heard, M.D.⁶
Pamela A. Lipsett, M.D.⁷
Henry Masur, M.D.¹
Leonard A. Mermel, D.O., Sc.M.⁸
Michele L. Pearson, M.D.⁹
Issam I. Raad, M.D.¹⁰
Adrienne Randolph, M.D., M.Sc.¹¹
Mark E. Rupp, M.D.¹²
Sanjay Saint, M.D., M.P.H.¹³

1National Institutes of Health, Bethesda, Maryland

2Infusion Nurses Society, Norwood, Massachusetts

3Greenwich Hospital, Greenwich, Connecticut

4University of Washington, Seattle, Washington

5Wheaton Franciscan Healthcare-St. Joseph, Milwaukee, Wisconsin

6 University of Massachusetts Medical School, Worcester, Massachusetts

7Johns Hopkins University School of Medicine, Baltimore, Maryland

8Warren Alpert Medical School of Brown University and Rhode Island Hospital, Providence, Rhode Island

9 Centers for Disease Control and Prevention, Atlanta, Georgia

10MD Anderson Cancer Center, Houston, Texas

11The Children's Hospital, Boston, Massachusetts

12University of Nebraska Medical Center, Omaha, Nebraska

13Ann Arbor VA Medical Center and University of Michigan, Ann Arbor, Michigan

36 Introduction

37

38 These guidelines have been developed for practitioners who insert catheters and

39 for persons responsible for surveillance and control of infections in hospital, outpatient,

40 and home healthcare settings. This report was prepared by a working group comprised of

Deleted: ing

41 members from professional organizations representing the disciplines of critical care

42 medicine, infectious diseases, healthcare infection control, surgery, anesthesiology,

43 interventional radiology, pulmonary medicine, pediatric medicine, and nursing. The

44 working group was led by the Society of Critical Care Medicine (SCCM), in

45 collaboration with the Infectious Disease Society of America (IDSA), Society for

46 Healthcare Epidemiology of America (SHEA), Surgical Infection Society (SIS),

47 American College of Chest Physicians (ACCP), American Thoracic Society (ATS),

48 American Society of Critical Care Anesthesiologists (ASCCA), Association for

49 Professionals in Infection Control and Epidemiology (APIC), Infusion Nurses Society

50 (INS), Oncology Nursing Society (ONS), Society of Cardiovascular and Interventional

51 Radiology (SCVIR), American Academy of Pediatrics (AAP), and the Healthcare

52 Infection Control Practices Advisory Committee (HICPAC) of the Centers for Disease

53 Control and Prevention (CDC) and is intended to replace the Guideline for Prevention of

54 Intravascular Device-Related Infections published in 2002. These guidelines are intended

55 to provide evidence-based recommendations for preventing catheter-related infections.

56 Major areas of emphasis include 1) educating and training healthcare personnel who

57 insert and maintain catheters; 2) using maximal sterile barrier precautions during central

58 venous catheter insertion; 3) using a 2% chlorhexidine preparation for skin antisepsis; 4)

59 avoiding routine replacement of central venous catheters as a strategy to prevent

60 infection; and 5) using antiseptic/antibiotic impregnated short-term central venous
61 catheters and chlorhexidine impregnated sponge dressings if the rate of infection is high
62 despite adherence to other strategies (i.e., education and training, maximal sterile barrier
63 precautions, and 2% chlorhexidine for skin antisepsis). These guidelines also emphasize
64 performance improvement by implementing bundled strategies, documenting and
65 reporting rates of [adherence to](#) all components of the bundle as benchmarks for
66 quality assurance and performance improvement.

Deleted: compliance rates with

67 As in previous guidelines issued by CDC and HICPAC, each recommendation is
68 categorized on the basis of existing scientific data, theoretical rationale, applicability, and
69 economic impact. The CDC/HICPAC system for categorizing recommendations is as
70 follows:

71 Category IA. Strongly recommended for implementation and strongly supported by well-
72 designed experimental, clinical, or epidemiologic studies.

73 Category IB. Strongly recommended for implementation and supported by some
74 experimental, clinical, or epidemiologic studies, and a strong theoretical rationale.

75 Category IC. Required by state or federal regulations, rules, or standards.

76 Category II. Suggested for implementation and supported by suggestive clinical or
77 epidemiologic studies or a theoretical rationale.

78 Unresolved issue. Represents an unresolved issue for which evidence is insufficient or no
79 consensus regarding efficacy exists.

80 Intravascular Catheter-Related Infections in Adult and Pediatric Patients: An

81 Overview

82 Background

83 In the United States, 15 million central vascular catheter (CVC) days (i.e., the
84 total number of days of exposure to CVCs by all patients in the selected population
85 during the selected time period) occur in intensive care units (ICUs) each year [1].
86 Catheter-related bloodstream infections (CRBSI) independently increase hospital costs
87 and length of stay [2-5], but have not been shown to independently increase mortality.
88 While 80,000 CVC-associated BSIs occur in ICUs each year [1], a total of 250,000 cases
89 of CVC-associated BSIs have been estimated to occur annually, if entire hospitals are
90 assessed [6]. By several analyses, the cost of CVC-associated BSI is substantial, both in
91 terms of morbidity and financial resources expended. To improve patient outcome and to
92 reduce healthcare costs, there is considerable interest by healthcare personnel, insurers,
93 regulators, and patient advocates reducing the incidence of these infections. This effort
94 should be multidisciplinary, involving healthcare personnel who order the insertion and
95 removal of CVCs, those personnel who insert and maintain intravascular catheters,
96 infection control personnel, healthcare managers from the CEO down to those who
97 allocate resources, and patients who are capable of assisting in the care of their catheters.
98 Personnel should recognize that the goal of an effective prevention program is continuous,
99 reduction in catheter-related infections. Elimination of catheter-related infection is a
100 laudable goal; demonstrating that elimination of CRBSIs can be sustained and encompass
101 CVCs placed at all points of care, e.g., ICU, med-surg, home care, etc., is challenging.
102 There are programs that have demonstrated success, but most programs will recognize
103 some catheter-related infections over time. The goal of the measures discussed in this
104 document is to reduce the rate to as low as feasible given the specific patient population

Comment [JDS1]: ? MRSA and VRE
CLABSIs associated with increased
mortality

Deleted: the

105 being served, the universal presence of microorganisms in the human environment, and
106 the limitations of current strategies and technologies.

107 Terminology and Estimates of Risk

108 The terminology used to identify different types of catheters is confusing, because
109 many clinicians and researchers use different aspects of the catheter for informal
110 reference. A catheter can be designated by the type of vessel it occupies (e.g., peripheral
111 venous, central venous, or arterial); its intended life span (e.g., temporary or short-term
112 versus permanent or long-term); its site of insertion (e.g., subclavian, femoral, internal
113 jugular, peripheral, and peripherally inserted central catheter [PICC]); its pathway from
114 skin to vessel (e.g., tunneled versus nontunneled); its physical length (e.g., long versus
115 short); or some special characteristic of the catheter (e.g., presence or absence of a cuff,
116 impregnation with heparin, antibiotics or antiseptics, and the number of lumens). To
117 accurately define a specific type of catheter, all of these aspects should be described
118 (Table 1).

Comment [JDS2]: definitions of short term, long term

119 The rate of all catheter-related infections, including local infections and systemic
120 infections, is difficult to determine. Potentially infectious episodes must be evaluated
121 clinically and microbiologically and documented in the record; the data must be reviewed
122 by well informed and fairly adjudicated personnel as to whether an episode is due to
123 infection or contamination and if infection is present, whether it is related to the CVC or
124 to a secondary source. Although CRBSI is a suitable parameter because it represents the
125 most serious form of catheter-related infection, it is often problematic to precisely
126 establish the diagnosis given the clinical setting of the patient (the catheter is not always
127 removed), limited availability of microbiologic methods (many labs do not use

Comment [JDS3]: not sure this is an accepted use of this word

Formatted: Highlight

128 quantitative blood cultures or differential time to positivity), and support by direct care
129 personnel (labeling must be accurate). Given these challenges, simpler automated
130 methods relying on microbiological data alone, albeit less precise, may offer convenient
131 alternatives to manual surveillance methods. Simplified objective criteria may be
132 potentially superior to clinical criteria in identifying the true differences in CRBSI rates
133 between institutions [7-10].

134 Healthcare personnel should recognize the difference between the surveillance
135 definition (i.e., the definition that is used to benchmark institutions reporting to the
136 National Healthcare Safety Network [NHSN] and clinical definitions. The NHSN
137 surveillance definition is for BSIs, including central-line associated BSIs, when other
138 documented sites of infection have been excluded
139 (http://www.cdc.gov/nhsn/PDFs/pscManual/4PSC_CLABSCurrent.pdf)[11]. That is, the
140 surveillance definition overestimates the true incidence of CRBSI because not all BSIs
141 originate from a catheter. Some bacteremias are secondary BSIs from unrecognized
142 sources (e.g., postoperative surgical sites, intra-abdominal infections, and hospital-
143 associated pneumonia or urinary tract infections). Thus, surveillance definitions are really
144 definitions for catheter-associated BSIs. Within this definition is opportunity for
145 subjective bias since some reviewers may be more prone than others to attribute the BSI
146 to other sources based on only vague, unconvincing information. This provides
147 opportunities for published rates of BSIs to be influenced by assessment teams with
148 motivation to record low rates.

149 A more rigorous definition might include only those BSIs for which other sources
150 were excluded by careful examination of the patient record, and where a culture of the

151 catheter-tip demonstrated substantial colonies of an organism identical to those found in
152 the bloodstream. Interpreting blood cultures drawn from catheters presents its own set of
153 challenges, but clinical definitions have been developed to take the results of such blood
154 cultures into account when establishing a diagnosis of CRBSI [12]. Such a clinical
155 /microbiological definition would focus on catheter-related BSIs. Therefore, to
156 accurately compare a healthcare facility's CRBSI infection rate to published data,
157 comparable definitions also should be used.

Comment [JDS4]: Important to include information on why catheter tips should not be cultured routinely.

158 CDC and The Joint Commission recommend that the rate of catheter-associated
159 BSIs (CABSI) be expressed as the number of CABSI per 1,000 CVC days [13, 14].
160 This parameter provides longitudinal data not expressed when the rate is expressed as the
161 number of catheter-associated infections per 100 catheters (or percentage of catheters
162 studied), because it accounts for BSIs over time and, therefore, adjusts risk for the
163 number of days the catheter is in use.

Comment [JDS5]: "because this denominator adjusts for duration of risk"

164 **Epidemiology and Microbiology in Adult and Pediatric Patients**

165 National estimates of CABSI rates are available through CDC's NHSN
166 (www.cdc.gov/nhsn). The most recently published NHSN data represent reports from
167 621 hospitals in 45 States and the District of Columbia that monitor infections in one or
168 more ICUs and/or non-ICUs (e.g., patient care areas, wards) [15]. Because BSI rates are
169 influenced by patient-related factors, such as severity of illness and type of illness (e.g.,
170 third-degree burns versus post-cardiac surgery), by catheter-related factors, (such as the
171 condition under which the catheter was placed and catheter type), and by institutional
172 factors (e.g., bed-size, academic affiliation), these aggregate, risk-adjusted rates can be

173 used as benchmarks against which hospitals can make intra- and inter-facility
174 comparisons.

175 Among hospitals participating in NHSN during 2006, the reported pooled mean
176 rates of central venous CABSIs ranged from 1.3/1000 catheter days on inpatient
177 medical/surgical wards to 5.6/1000 catheter days in burn ICUs (Table 2). In neonatal
178 nurseries for infants weighing less than 1,000 grams (Level III), central line-associated
179 BSI rates ranged from 3.3-3.7/1,000 catheter days [15]. In these nurseries, umbilical
180 catheter rates also varied by birth weight category, ranging from 0.9/1,000 catheter days
181 among neonates weighing above 1,500 grams to 4.7/1,000 catheter days among neonates
182 weighing 750 grams or less [15]. Secular trends suggest a reduction in the incidence of
183 central venous CABSIs occurring in ICUs during the past 20 years.

Comment [JDS6]: Would use CLABSI instead of CABSIs because that is what is used by NHSN. It is too confusing to go back and forth between the two.

Deleted: over

184 The most commonly reported causative pathogens for hospital acquired BSIs
185 remain coagulase-negative staphylococci, *Staphylococcus aureus*, enterococci, and
186 *Candida* spp. [16]. Gram negative bacilli accounted for 19% and 21% of catheter
187 associated BSIs reported to CDC [17] and the Surveillance and Control of Pathogens of
188 Epidemiological Importance (SCOPE) database, respectively [16].

189 For all common pathogens causing CRBSIs, antimicrobial resistance is a problem,
190 particularly in ICUs. Although methicillin-resistant *Staphylococcus aureus* (MRSA) now
191 accounts for more than 50% of all *Staphylococcus aureus* isolates obtained in ICUs, the
192 incidence of MRSA CABSIs has decreased in recent years, most likely, as a result of

Deleted: perhaps

193 prevention efforts [18] For gram negative rods, antimicrobial resistance to third
194 generation cephalosporins among *Klebsiella pneumoniae* and *E. coli* have increased

195 significantly as did imipenem and ceftazidime resistance among *Pseudomonas aeruginosa*

196[17]. *Candida* spp. are noted to be increasingly resistant to fluconazole,

Deleted: increasingly

Deleted: resistant

197 As in adults, the majority of BSIs in children are associated with the use of an

198 intravascular catheter. From 2002 through 2004, the pooled mean CABS rate for all

199 pediatric ICUs reporting data to NNIS was 6.6 per 1,000 catheter days [14]. This rate has

200 decreased compared to the 1995-2000 data, but is consistently higher than that reported in

201 adult medical-surgical ICUs during the 2002-2004 time period. Umbilical catheter and CVC-
associated BSI rates for

202 neonatal ICUs ranged from 3.7-4.7 per 1,000 catheter days in children with birth weight

203 <750 gram to 1.0-2.0 per 1,000 catheter days in children whose birth weight was >2,500

204 gram [19]. Catheter utilization ratios were comparable in adult and pediatric ICUs [20,

Deleted: e

205 21].

206 The distribution of types of organisms causing infection is similar in pediatric

207 ICUs and adult ICUs [21]. As in adults, the majority of CRBSIs in children are caused

208 by coagulase-negative staphylococci. During 1992-1999, these bacteria accounted for

209 37.7% of BSIs in pediatric ICUs reporting to NNIS [21]. Among neonates with

210 percutaneously placed central venous catheters, coagulase-negative staphylococci are

211 responsible for 75% of CRBSIs [22, 23].

212 Pathogenesis

213 There are four recognized routes for contamination of catheters: 1) migration of

214 skin organisms at the insertion site into the cutaneous catheter tract and along the surface

215 of the catheter with colonization of the catheter tip; this is the most common route of

216 infection for short-term catheters [24-26]; 2) direct contamination of the catheter or

217 catheter hub by contact with hands or contaminated fluids or devices [27, 28]; 3) less

218 commonly, catheters might become hematogenously seeded from another focus of
219 infection [29]; and 4) rarely, infusate contamination might lead to CRBSI [30].

220 Important pathogenic determinants of catheter-related infection are 1) the material
221 of which the device is made; 2) the host factors consisting of protein adhesions, such as
222 fibrin and fibronectin that form a sheath around the catheter [31]; and 3) the intrinsic
223 virulence factors of the infecting organism, including the extracellular polymeric
224 substance (EPS) produced by the adherent organisms [32]. Some catheter materials also
225 have surface irregularities that enhance the microbial adherence of certain species (e.g., *S.*
226 *epidermidis* and *C. albicans*) [33, 34]. Catheters made of these materials are especially
227 vulnerable to microbial colonization and subsequent infection. After the formation of the
228 fibrin sheath, silastic catheters are more prone to catheter infections than polyurethane
229 catheters [31]. On the other hand, biofilm formation by *C. albicans* occurs more
230 intensely on silicone elastomer catheter surfaces than polyurethane catheters [33].

Comment [JDS7]: Wouldn't immune system status be included as a host factor?

231 Modification of the biomaterial surface properties has been shown to influence the ability
232 of *C. albicans* to form biofilm [34]. Additionally, certain catheter materials are more
233 thrombogenic than others, a characteristic that also might predispose to catheter
234 colonization and catheter-related infection [35, 36]. This association has led to emphasis
235 on preventing catheter-related thrombus as an additional mechanism for reducing CRBSI
236 [37, 38].

Comment [JDS8]: Not really "on the other hand"; additionally might be a better word.

237 The adherence properties of a given microorganism in relationship to host factors
238 are also important in the pathogenesis of catheter-related infection. For example, *S.*
239 *aureus* can adhere to host proteins (e.g., fibrinogen, fibronectin) commonly present on
240 catheters by expressing clumping factors (ClfA and ClfB) that bind to the protein

241 adhesins [31, 36, 39, 40]. Furthermore, adherence is enhanced through the production of
242 EPS by microbial organisms, such as coagulase-negative staphylococci [41, 42], *S.*
243 *aureus* [43], *Pseudomonas aeruginosa* [44], and *Candida* spp. [45], consisting mostly of
244 an exopolysaccharide that forms a microbial biofilm layer [32, 46]. This biofilm matrix is
245 enriched by divalent metallic cations, such as calcium, magnesium and iron, which make
246 it a solid enclave for microbial organisms to embed themselves [47-49]. In the presence
247 of catheters, this biofilm potentiates the pathogenicity of various microbes by allowing
248 them to withstand host defense mechanisms (e.g., acting as a barrier to engulfment and
249 killing by polymorphonuclear leukocytes) or by making them less susceptible to
250 antimicrobial agents (e.g., forming a matrix that binds antimicrobials before their contact
251 with the organism cell wall) [42, 50, 51]. Some *Candida* spp., in the presence of glucose-
252 containing fluids, produce slime similar to that of their bacterial counterparts, potentially
253 explaining the increased proportion of BSIs caused by fungal pathogens among patients
254 receiving parenteral nutrition fluids [52].

Formatted: Portuguese (Brazil)

Comment [JDS9]: Is there anything that can be done to alter the biofilms as a prevention strategy?

255 Strategies for Prevention of Catheter-Related Infections in Adult and Pediatric 256 Patients

257 Education, training and staffing

258 Recommendations

Comment [JDS10]: Should there be a recommendation for joining collaboratives to reduce CLABSI rates?

- 259 1. Educate healthcare personnel regarding the indications for intravascular catheter use,
260 proper procedures for the insertion and maintenance of intravascular catheters, and
261 appropriate infection control measures to prevent intravascular catheter-related infections
262 [53-61]. Category IA

263 2. Periodically assess knowledge of and adherence to guidelines for all persons who are
264 involved in the insertion and maintenance of intravascular catheters [53-61]. Category IA
265 3. Designate only trained personnel who demonstrate competence for the insertion and
266 maintenance of peripheral and central intravascular catheters. [60-74]. Category IA
267 4. Ensure appropriate nursing staff levels in ICUs to minimize the incidence of catheter-
268 related BSIs. Observational studies suggest a ratio of 2:1 in ICUs where nurses are
269 managing patients with CVCs [75-77]. Category IB

270 **Background**

271 Well-organized programs that enable healthcare personnel to become educated
272 and to provide, monitor, and evaluate care are critical to the success of this effort. Reports
273 spanning the past four decades have consistently demonstrated that risk for infection
274 declines following standardization of aseptic care [53, 58, 60, 61, 78-80] and that
275 insertion and maintenance of intravascular catheters by inexperienced staff
276 increase the risk for catheter colonization and CRBSI [61, 81]. Specialized "IV teams"
277 have shown unequivocal effectiveness in reducing the incidence of catheter-related
278 infections, associated complications, and costs [62-72]. Additionally, infection risk
279 increases with nursing staff reductions below a critical level [76].

280 **Site selection**

281 **Recommendations for peripheral catheters and midline catheters**

282 1. In adults, use an upper-extremity site for catheter insertion. Replace a catheter inserted
283 in a lower extremity site to an upper extremity site as soon as possible [82, 83]. Category
284 IB

Deleted: might

- 285 2. In pediatric patients, the upper or lower extremities or the scalp can be used as the
286 catheter insertion site [82, 83]. Category II
- 287 3. Select catheters on the basis of the intended purpose and duration of use, known
288 infectious and non-infectious complications (e.g., phlebitis and infiltration), and
289 experience of individual catheter operators [83-85]. Category IB
- 290 4. Avoid the use of steel needles for the administration of fluids and medication that
291 might cause tissue necrosis, if extravasation occurs [83-85]. Category IA
- 292 5. Use a midline catheter or peripherally inserted central catheter (PICC), instead of a
293 short peripheral catheter, when the duration of IV therapy will likely exceed six days [83-
294 85]. Category IB
- 295 **Recommendations for central venous catheters**
- 296 6. Weigh the risk and benefits of placing a central venous device at a recommended site
297 to reduce infectious complications against the risk for mechanical complications (e.g.,
298 pneumothorax, subclavian artery puncture, subclavian vein laceration, subclavian vein
299 stenosis, hemothorax, thrombosis, air embolism, and catheter misplacement) [25, 86-
300 101]. Category IA
- 301 7. Use a subclavian site, rather than a jugular or a femoral site, in adult patients to
302 minimize infection risk for nontunneled CVC placement [25, 99, 100]. Category IA
- 303 8. No recommendation can be made for a preferred site of insertion to minimize infection
304 risk for a tunneled CVC. Unresolved issue
- 305 9. Place catheters used for hemodialysis and pheresis in a jugular or femoral vein, rather
306 than a subclavian vein, to avoid venous stenosis [101-105]. Category IA

307 10. Use ultrasound guidance to place central venous catheters to reduce the number of
308 cannulation attempts and mechanical complications if this technology is available [106,
309 107]. Category 1B

Comment [JDS11]: Should you include use of a sterile sleeve on the ultrasound probe and cord?

310 11. Promptly remove any intravascular catheter that is no longer essential [108, 109].

311 Category IA

312 Background

313 The site at which a catheter is placed influences the subsequent risk for catheter-
314 related infection and phlebitis. The influence of site on the risk for catheter infections is
315 related in part to the risk for thrombophlebitis, density of local skin flora, and risk of
contamination by infectious body fluids, e.g., stool, saliva.

Deleted: and

316 Phlebitis has long been recognized as a risk for infection. For adults, lower
317 extremity insertion sites are associated with a higher risk for infection than are upper
318 extremity sites [110-112]. In addition, hand veins have a lower risk for phlebitis than do
319 veins on the wrist or upper arm [113]. As in adults, the use of peripheral venous catheters
320 in pediatric patients might be complicated by phlebitis, infusion extravasation, and
321 catheter infection [114]. Catheter location, infusion of parenteral nutritional fluids with
322 continuous IV lipid emulsions, and length of ICU stay before catheter insertion have all
323 increased pediatric patients' risk for phlebitis. However, contrary to the risk in adults, the
324 risk for phlebitis in children has not increased with the duration of catheterization [114,
325 115].

326 The density of skin flora at the catheter insertion site is a major risk factor for
327 CRBSI. Authorities recommend that CVCs be placed in a subclavian site, instead of a
328 jugular or femoral site, to reduce the risk for infection. No single trial has satisfactorily
329 compared infection rates for catheters placed in jugular, subclavian, and femoral vein. In

330 retrospective observational studies, catheters inserted into an internal jugular vein have
331 usually been associated with higher risk for colonization and/or CRBSI than those
332 inserted into a subclavian or femoral vein [25, 86-95]. Similar findings were noted in
333 neonates in a single retrospective study [116].

334 Femoral catheters have been demonstrated to have high colonization rates
335 compared to subclavian and internal jugular sites when used in adults and, in some
336 studies, higher rates of CRBSIs [88, 93-95, 98, 99, 117]. Femoral catheters should also be
337 avoided, when possible, because they are associated with a higher risk for deep venous
338 thrombosis than are internal jugular or subclavian catheters [96-98, 101, 118]. One study
339 [86] found that the risk of infection associated with catheters placed in the femoral vein is
340 accentuated in obese patients. In contrast to adults, studies in pediatric patients have
341 demonstrated that femoral catheters have a low incidence of mechanical complications
342 and might have an equivalent infection rate to that of nonfemoral catheters [119-122].
However, diapered children may have an increased risk of contamination with stool.

343 Thus, in adult patients, a subclavian site is preferred for infection control purposes,
344 although other factors (e.g., the potential for mechanical complications, risk for
345 subclavian vein stenosis, and catheter-operator skill) should be considered when choosing a
catheter insertion site.

Deleted: deciding
346 . where to place the catheter.

347 In two meta-analyses, the use of dynamic two-dimensional ultrasound for the
348 placement of CVCs substantially decreased mechanical complications and reduced the
349 number of attempts at required cannulation and failed attempts at cannulation compared
350 with the standard landmark placement [106, 107]. Evidence favors the use of two
351 dimensional ultrasound guidance over Doppler ultrasound guidance [106]. Site selection
352 should be guided by patient comfort, ability to secure the catheter, and maintenance of

353 asepsis as well as patient-specific factors (e.g., preexisting catheters, anatomic deformity,
354 and bleeding diathesis), relative risk of mechanical complications (e.g., bleeding and
355 pneumothorax), the availability of bedside ultrasound, the experience of the person
356 inserting the catheter, and the risk for infection.

357 Catheters should be inserted as great a distance as possible from open wounds [and from](#)
358 [ostomy sites](#). In

358 one study, catheters inserted close to open burn wounds were 1.79 times more likely to be
359 colonized and 5.12 times more likely to be associated with bacteremia than catheters
360 inserted farther from the wounds [123].

361 **Type of Catheter Material**

362 Polytetrafluoroethylene or polyurethane catheters have been associated with fewer
363 infectious complications than catheters made of polyvinyl chloride or polyethylene [124-
364 126]. Steel needles used as an alternative to catheters for peripheral venous access have
365 the same rate of infectious complications as do polytetrafluoroethylene catheters [83, 84].
366 However, the use of steel needles frequently is complicated by infiltration of intravenous
367 (IV) fluids into the subcutaneous tissues, a potentially serious complication if the infused
368 fluid is a vesicant [84].

369 **Hand Hygiene and Aseptic Technique**

370 **Recommendations**

371 1. Perform hand hygiene procedures, either by washing hands with
372 antiseptic containing soap and water or with waterless alcohol-based hand rubs (ABHR).
373 Hand hygiene should be performed before and after palpating catheter insertion sites as
374 well as before and after inserting, replacing, accessing, repairing, or dressing an
375 intravascular catheter. Palpation of the insertion site should not be performed after the

Deleted: conventional

376 application of antiseptic, unless aseptic technique is maintained [58, 127-131]. Category
377 IA

378 2. Maintain aseptic technique for the insertion and care of intravascular catheters [25,
379 132-134]. Category IA

380 3. Wear clean gloves, rather than sterile gloves, for the insertion of peripheral
381 intravascular catheters, if the access site is not touched after the application of skin
382 antiseptics. Category IC

383 4. Wear sterile gloves for the insertion of arterial, central, and midline
384 catheters [25, 132-134]; and change these gloves, if a catheter is being

Deleted: S

Deleted: should be worn

Deleted: should be changed

385 exchanged over a guidewire (thereby contaminating the gloves) and a new sterile catheter
386 is then handled. Category IA

387 4. Wear either clean or sterile gloves when changing the dressing on intravascular
388 catheters. Category IC

389 **Background**

390 Hand hygiene before catheter insertion or maintenance, combined with proper
391 aseptic technique during catheter manipulation, provides protection against infection
392 [58]. Proper hand hygiene can be achieved through the use of either a waterless, alcohol-
393 based product [135] or an antibacterial soap and water with adequate rinsing [127].
394 Appropriate aseptic technique does not necessarily require sterile gloves for insertion of
395 peripheral catheters; a new pair of disposable nonsterile gloves can be used in
396 conjunction with a "no-touch" technique for the insertion of peripheral venous catheters.
397 Sterile gloves must be worn for placement of central catheters since a "no-touch"
398 technique is not possible.

399 Maximal Sterile Barrier Precautions

400 Recommendations

401 1. Use maximal sterile barrier precautions, including the use of a cap, mask, sterile gown,
402 sterile gloves, and a large sterile full body drape, for the insertion of CVCs, PICCs, or
403 guidewire exchange [60, 132, 136, 137]. Category IB

404 2. Use a sterile sleeve to protect pulmonary artery catheters during insertion [138].

405 Category IB

406 Background

407 Maximum sterile barrier (MSB) precautions is defined as wearing a sterile gown,
408 sterile gloves, and cap and using a full body drape (similar to the drapes used in the
409 operating room) during the placement of CVC. Maximal sterile barrier precautions during
410 insertion of CVC were compared with sterile gloves and a small drape in a randomized
411 controlled trial. The MSB group had fewer episodes of both catheter colonization (RR =
412 0.32, 95% CI, 0.10-0.96, P = .04) and CR-BSI (RR = 0.16, 95% CI, 0.02-1.30, P = .06).

Formatted: French (France)

413 In addition, the group with MSB had infections that occurred much later and contained
414 gram negative, rather than gram positive, organisms [132]. A study designed to examine
415 pulmonary artery catheters also secondarily demonstrated that use of MSB precautions
416 was one of the items that lowered risk of infection [25]. Another study evaluated an
417 educational program directed at improving infection control practices, especially MSB. In
418 this study, MSB use increased and CRBSI decreased [60]. A small trial demonstrated an
419 reduced risk of skin colonization at the insertion site with maximal barrier precautions
420 [OR 3.40, 95%CI 1.32 to 3.67] [136].

421 Skin Preparation

422 Recommendations

423

424 1. Prepare clean skin with 70% alcohol before peripheral venous catheter insertion [139].

425 Category IA

426 2. Prepare clean skin site with a 2% chlorhexidine-based preparation before central

427 venous catheter insertion and during dressing changes. If there is a contraindication to

428 chlorhexidine, tincture of iodine, an iodophor, or 70% alcohol can be used as alternatives

429 [140, 141]. Category IA

430 3. No recommendation can be made for the safety or efficacy of chlorhexidine in infants

431 aged <2 months. Unresolved issue

Comment [JDS12]: Suggest including in the discussion experience with CHG in neonates. It is being used extensively and there are some data.

432 4. Allow povidone iodine to remain on the skin for at least 2 minutes or longer for the

433 antibacterial properties to take effect, if it is not yet dry before catheter insertion. The

434 antibacterial properties of chlorhexidine work on contact, and chlorhexidine does not

435 require a minimum 2- minute drying time before proceeding. Catheter insertion may

Comment [JDS13]: So, no drying time at all is required with CHG?

436 begin as soon as the chlorhexidine is dry[140, 141]. Category IB

437 Background

438

439 Two well-designed studies evaluating the chlorhexidine-containing cutaneous

440 antiseptic regimen in comparison with either povidone iodine or alcohol for the care of an

441 intravascular catheter insertion site have shown lower rates of catheter colonization or

442 CRBSI associated with the chlorhexidine preparation [140, 141]. When 0.5% tincture of

443 chlorhexidine was compared with 10% povidone iodine, no differences were seen in

444 CVC colonization or in CRSBI [142]. In a three-armed study (2% aqueous chlorhexidine

445 gluconate vs 10% povidone-iodine vs 70% alcohol), 2% aqueous chlorhexidine gluconate

446 tended to decrease CRBSI compared with 10% povidone iodine or 70% alcohol [140]. A
447 meta-analysis of 4,143 catheters suggested that chlorhexidine preparation, rather than
448 povidone iodine, reduced the risk of catheter-related infection by 49% (95% CI 0.28 to
449 0.88) [143]. An economic decision analysis based on available evidence suggested that
450 the use of chlorhexidine, rather than povidone iodine, for CVC care would result in a
451 1.6% decrease in the incidence of CRBSI, a 0.23% decrease in the incidence of death,
452 and a savings of \$113 per catheter used [144]. While 2% chlorhexidine has become a
453 standard antiseptic for skin preparation for the insertion of both central and peripheral
454 venous catheters, 5% povidone iodine solution in 70% ethanol was associated with a
455 substantial reduction of CVC-related colonization and infection compared with 10%
456 aqueous povidone iodine [145].

Comment [JDS14]: This implies that there is an increased risk of death associated with CLABSI, whereas, earlier in the document, it said that such was not the case.

457 **Catheter site dressing regimens**

458 **Recommendations**

- 459 1. Use either sterile gauze or sterile, transparent, semi-permeable dressing to cover the
460 catheter site [146-149]. Category IA
- 461 2. If the patient is diaphoretic or if the site is bleeding or oozing, use gauze dressing until
462 this is resolved [146-149]. Category II
- 463 3. Replace catheter site dressing if the dressing becomes damp, loosened, or visibly soiled
464 [146, 147]. Category IB
- 465 4. Do not use topical antibiotic ointment or creams on insertion sites, except for dialysis
466 catheters, because of their potential to promote fungal infections and antimicrobial
467 resistance [150, 151]. Category IB

Comment [JDS15]: What about compatibility of the creams/ointments with the hemodialysis catheters?

468 5. Do not submerge the catheter or catheter site in water. Showering should be permitted
469 [only](#) if precautions can be taken to reduce the likelihood of introducing organisms into the
470 catheter (e.g., if the catheter and connecting device are protected with an impermeable
471 cover during the shower) [152, 153]. Category II

Comment [JDS16]: No data for IB?

472 6. Replace dressings used on short-term CVC sites every 2 days for gauze dressings and
473 at least every 7 days for transparent dressings, except in those pediatric patients in which
474 the risk for dislodging the catheter may outweigh the benefit of changing the dressing
475 [149]. Category IB

476 7. Replace dressings used on tunneled or implanted CVC sites no more than once per
477 week, until the insertion site has healed [149]Category IB

478 8. No recommendation can be made regarding the necessity for any dressing on well-
479 healed exit sites of long-term cuffed and tunneled CVCs. Unresolved issue

480 9. Ensure that catheter site care is compatible with the catheter material [154, 155].

481 Category IB

482 10. Use a sterile sleeve for all pulmonary artery catheters [138]. Category IB

483 11. Use a chlorhexidine-impregnated sponge dressing for temporary short-term catheters

484 in patients older than 2 months of age, if the CRBSI rate is higher than the institutional

Comment [JDS17]: More specific info. on age, BW criteria?

485 goal, despite adherence to basic CRBSI prevention measures, including education and

486 training, use of chlorhexidine for skin antisepsis, and MSB [22, 156-158]. Category 1B

487 **Background**

488 Transparent, semi-permeable polyurethane dressings permit continuous visual
489 inspection of the catheter site and require less frequent changes than do standard gauze
490 and tape dressings. In the largest controlled trial of dressing regimens on peripheral

491 catheters, the infectious morbidity associated with the use of transparent dressings on
492 approximately 2,000 peripheral catheters was examined [126]. Data from this study
493 suggest that the rate of colonization among catheters dressed with transparent dressings
494 (5.7%) is comparable to that of those dressed with gauze (4.6%) and that no clinically
495 substantial differences exist in either the incidences of catheter site colonization or
496 phlebitis. Furthermore, these data suggest that transparent dressings can be safely left on
497 peripheral venous catheters for the duration of catheter insertion without increasing the
498 risk for thrombophlebitis [126].

499 A meta-analysis has assessed studies that compared the risk for CRBSIs for
500 groups using transparent dressings versus groups using gauze dressing [159]. The risk for
501 CRBSIs did not differ between the groups. The choice of dressing can be personal
502 preference. If blood is oozing from the catheter insertion site, gauze dressing is preferred.
503 Another systemic review of randomized controlled trials comparing gauze and tape to
504 transparent dressings found no significant differences in CRBSIs, catheter tip
505 colonization, or skin colonization between dressing types [160].

Formatted: Highlight

Deleted: a

Deleted: matter of

506 Chlorhexidine impregnated dressings have been used to reduce the risk of CRBSI.
507 In the largest multicenter randomized controlled trial published to date comparing
508 chlorhexidine impregnated sponge dressings vs standard dressings in ICU patients, rates
509 of CRIs were reduced even when background rates of infection were low. In this study,
510 1636 patients (3778 catheters, 28 931 catheter-days) were evaluated. The chlorhexidine-
511 impregnated dressings decreased the rates of major CRIs (10/1953 [0.5%], 0.6 per 1000
512 catheter-days vs 19/1825 [1.1%], 1.4 per 1000 catheter-days; hazard ratio [HR], 0.39
513 [95% confidence interval {CI}, 0.17-0.93]; $P = .03$) and CRBSIs (6/1953 catheters, 0.40

Formatted: Highlight

514 per 1000 catheter-days vs 17/1825 catheters, 1.3 per 1000 catheter-days; HR, 0.24 [95%
515 CI, 0.09-0.65]) [156]. A randomized controlled study of 140 children used polyurethane
516 or a chlorhexidine impregnated dressing showed no statistical difference in BSIs;
517 however, the chlorhexidine group had lower rates of CVC colonization [158]. In 601
518 cancer patients receiving chemotherapy, the incidence of CRBSI was reduced in patients
519 receiving the chlorhexidine sponge dressing compared to standard dressings (p=0.016,
520 relative risk 0.54; confidence interval 0.31-0.94) [161]. A meta-analysis that included
521 eight randomized controlled trials demonstrated that chlorhexidine impregnated sponges
522 are associated with a reduction of vascular and epidural catheter exit site colonization
523 (14.8% versus 26.9%, OR 0.47, 95% CI: 0.34 to 0.65) (overall 14.3% versus 27.2%, OR
524 0.40, 95% CI: 0.26–0.61; P < 0.0001), but no significant reduction in CRBSI (2.2%
525 versus 3.8%, OR 0.58, 95% CI: 0.29–1.14, P = 0.11) [157].

526 Although data regarding the use of a chlorhexidine impregnated sponge in
527 children are limited, one randomized, controlled study involving 705 neonates reported a
528 substantial decrease in colonized catheters in infants in the chlorhexidine sponge group
529 compared with the group that had standard dressings (15% versus 24%; RR = 0.6; 95%
530 CI = 0.5–0.9), but no difference in the rates of CRBSI or BSI without a source.

531 Chlorhexidine impregnated sponges were associated with localized contact dermatitis in
532 infants of very low birth weight. In 98 neonates with very low birth weight, 15 (15%)
533 developed localized contact dermatitis; four (1.5%) of 237 neonates weighing >1,000 g
534 developed this reaction (p < 0.0001). Infants with gestational age <26 weeks who had
535 CVCs placed at age <8 days were at increased risk for having localized contact
536 dermatitis, whereas no infants in the control group developed this local reaction [22].

537 Patient Cleansing

538 Recommendation

539 Use a 2% chlorhexidine wash daily to reduce CRBSI [162]. Category II

Comment [JDS18]: For adults only?

540 Background

541 Daily cleansing of ICU patients with a 2% chlorhexidine impregnated washcloth may be
542 a simple, effective strategy to decrease the rate of primary BSIs. In a single center study
543 of 836 ICU patients, patients receiving the chlorhexidine intervention were significantly
544 less likely to acquire a primary BSI (4.1 vs 10.4 infections per 1000 patient days;
545 incidence difference, 6.3 [95% confidence interval, 1.2-11.0) than those bathed with soap
546 and water [162].

547 Catheter Securement Devices

548 Recommendation

549 Use a sutureless securement device to reduce the risk of infection for PICCs [163].

550 Category II

551 Background

552 Catheter stabilization is recognized as an intervention to decrease the risk for
553 phlebitis, catheter migration and dislodgement, and may be advantageous in preventing
554 CRBSIs. Pathogenesis of CRBSI occurs via migration of skin flora through the
555 percutaneous entry site. Sutureless securement devices avoid disruption around the
556 catheter entry site and may decrease the degree of bacterial colonization. [163]. Using a
557 sutureless securement device also mitigates the risk of sharps injury to the healthcare
558 personnel from inadvertent needlestick injury.

Deleted: u

559 Antimicrobial/Antiseptic Impregnated Catheters and Cuffs

560 Recommendation

561 Use a chlorhexidine/silver sulfadiazine or minocycline/rifampin -impregnated CVC in
562 adults whose catheter is expected to remain in place >5 days if, after successful
563 implementation of a comprehensive strategy to reduce rates of CRBSI, the CRBSI rate
564 remains above the goal set by the individual institution based on benchmark rates (Tables
565 2 and 3) and local factors. The comprehensive strategy should include at least the
566 following three components: educating persons who insert and maintain catheters, use of
567 maximal sterile barrier precautions, and a 2% chlorhexidine preparation for skin
568 antiseptis during CVC insertion. Category IA

Comment [jds19]: ? and less than 30 days

569 Background

570 Certain catheters and cuffs that are coated or impregnated with antimicrobial or
571 antiseptic agents can decrease the risk for CRBSI and potentially decrease hospital costs
572 associated with treating CRBSIs, despite the additional acquisition cost of an
573 antimicrobial/antiseptic impregnated catheter [164]. Nearly all of the studies involving
574 antimicrobial/antiseptic-impregnated catheters have been conducted using triple-lumen,
575 uncuffed catheters in adult patients whose catheters remained in place <30 days. Most of
576 the studies have been conducted in adults; however, these catheters have been approved
577 by FDA for use in patients weighing >3 kg. Two non-randomized studies [165, 166] in
578 pediatric ICU patients suggest that these catheters may reduce risk of catheter-associated
579 infection. No antiseptic or antimicrobial impregnated catheters currently are available for
580 use in infants weighing <3 kg.

581 **Chlorhexidine/Silver sulfadiazine.**

582 Catheters coated with chlorhexidine/silver sulfadiazine only on the external
583 luminal surface have been studied as a means to reduce CRBSI. Two meta-analyses of
584 first-generation catheters [1, 167] demonstrated that such catheters reduced the risk for
585 CRBSI compared with standard non-coated catheters. The duration of catheter placement
586 in one study ranged from 5.1 to 11.2 days [168]. A second-generation catheter is now
587 available with chlorhexidine coating the internal surface extending into the extension set
588 and hubs while the external luminal surface is coated with chlorhexidine and silver
589 sulfadiazine. The external surface has three times the amount of chlorhexidine and
590 extended release of the surface bound antiseptics than that in the first generation
591 catheters. All three prospective, randomized studies of second-generation catheters
592 demonstrated a significant reduction in catheter colonization, but they were
593 underpowered to show a difference in CRBSI [169-171]. Prolonged anti-infective activity
594 provides improved efficacy in preventing infections [172]. Although rare, anaphylaxis
595 with the use of these chlorhexidine/silver sulfadiazine catheters has been observed [173-
596 176].

597 Chlorhexidine/silver sulfadiazine catheters are more expensive than standard
598 catheters. However, one analysis has suggested that the use of chlorhexidine/silver
599 sulfadiazine catheters should lead to a cost savings of \$68 to \$391 per catheter [177] in
600 settings in which the risk for CRBSI is high, despite adherence to other preventive
601 strategies (e.g., maximal barrier precautions and aseptic techniques). Use of these
602 catheters might be cost effective in ICU patients, burn patients, neutropenic patients, and
603 other patient populations in which the rate of infection exceeds 3.3 per 1,000 catheter
604 days [168].

Comment [jds20]: How was 3.3/1,000 derived? I think this is outdated with the more recent lower rates associated with bundled practices.

605 **Minocycline/Rifampin.**

606 In a multicenter randomized trial, CVCs impregnated on both the external and
607 internal surfaces with minocycline/rifampin were associated with lower rates of CRBSI
608 when compared with the first generation chlorhexidine/silver sulfadiazine impregnated
609 catheters [178]. The beneficial effect began after day 6 of catheterization. Silicone
610 minocycline/rifampin impregnated CVCs with an average dwell time of over 60 days
611 have been shown to be effective in reducing CRBSI [179]. No minocycline/rifampin-
612 resistant organisms were reported. Two trials demonstrated that use of these catheters
613 significantly reduced CRBSI compared to uncoated catheters [164, 179]. No
614 comparative studies have been published using the second-generation chlorhexidine/
615 silver sulfadiazine catheter. Although there have been concerns related to the potential for
616 development of resistance, several prospective clinical studies have shown that the risk is
617 low [180, 181]. Further, no resistance to minocycline or rifampin related to the use of the
618 catheter has been documented in the clinical setting. Two studies using decision model
619 analysis revealed these catheters were associated with superior cost savings compared
620 with first generation chlorhexidine/silver sulfadiazine catheters [182, 183]. Such analysis
621 needs to be done compared to the second-generation catheters. However, as baseline rates
622 of infection decrease and the cost of catheters decreases, the cost-benefit ratio will likely
623 change.

624 The decision to use chlorhexidine/silver sulfadiazine or minocycline/rifampin
625 impregnated catheters should be based on the need to enhance prevention of CRBSI after
626 bundled standard procedures have been implemented and shown to be practiced
627 consistently by auditing (e.g., educating personnel, using
627 maximal sterile barrier precautions, and using 2% chlorhexidine skin antisepsis) and then

628 balanced against the concern for emergence of resistant pathogens and the cost of
629 implementing this strategy.

630 **Platinum/Silver**

631 A combination platinum/silver impregnated catheter (i.e., a silver iontophoretic
632 catheter) is available for use in the United States. Several prospective, randomized studies
633 have been published comparing these catheters to uncoated catheters [184-187]. One
634 study showed a reduction in the incidence density of catheter colonization and CRBSI
635 [186], but the other studies found no difference in catheter colonization or CRBSI
636 between the impregnated catheter and a non-impregnated catheter [87, 184, 185].

637 **Systemic Antibiotic Prophylaxis**

638 **Recommendation**

639 Do not administer systemic antimicrobial prophylaxis routinely before insertion or during
640 use of an intravascular catheter to prevent catheter colonization or CRBSI [188].

641 Category IA

642 **Background**

643 Several studies have examined the role of systemic antibiotic prophylaxis in
644 prevention of catheter-related infection. A recent meta-analysis reviewed these studies in
645 oncology patients [188]. Four studies utilized a prophylactic glycopeptide prior to
646 catheter insertion. However, heterogeneity in these studies precludes any conclusion from
647 being reached about efficacy

648 In a study examining the effect of ongoing oral prophylaxis with rifampin and
649 novobiocin on catheter-related infection in cancer patients treated with interleukin-2
650 [189], a reduction in CRBSI was observed, even though 9 of 26 subjects (35%)

651 discontinued the prophylactic antibiotics due to side effects or toxicity. In non-oncology
652 patients, no benefit was associated with vancomycin administration prior to catheter
653 insertion in 55 patients undergoing catheterization for parenteral nutrition [190].
654 Similarly, extending perioperative prophylactic antibiotics in cardiovascular surgery
655 patients did not reduce central venous catheter colonization [191]. A recent Cochrane
656 review of prophylactic antibiotics in neonates with umbilical venous catheters concluded
657 that there is insufficient evidence from randomized trials to support or refute the use of
658 prophylactic antibiotics [192].

659 Late onset neonatal sepsis is often due to coagulase negative staphylococci and is
660 thought to frequently stem from infected central venous catheters. Five trials involved a
661 total of 371 neonates treated with vancomycin, either by continuous infusion via
662 parenteral nutrition or intermittent dosing or placebo. The infants treated with
663 vancomycin experienced less nosocomial sepsis (RR 0.11; 95% CI 0.05-0.24) and less
664 sepsis due to coagulase-negative staphylococci (RR 0.33; 95% CI 0.19-0.59) [193].
665 However, mortality and length of stay were not significantly different between the two
666 groups. There were insufficient data to evaluate the risk of development of vancomycin-
667 resistant organisms.

668 Antibiotic/Antiseptic Ointments

669 Recommendation

670 Use povidone iodine antiseptic ointment or bacitracin/neomycin/polymyxin B ointment at
671 the hemodialysis catheter exit site after catheter insertion and at the end of each dialysis
672 session only if this ointment does not interact with the material of the hemodialysis
673 catheter per manufacturer's recommendation [139, 194-198]. Category IB

674 Background

675 A variety of topical antibiotic or antiseptic ointments have been utilized in
676 attempts to lower the antimicrobial burden at the catheter insertion site and thus prevent
677 infection. A number of older studies, examining primarily peripheral venous catheters,
678 yielded varying conclusions [139, 199, 200]. In addition, the use of antibiotic ointments
679 that have limited antifungal activity may serve to increase colonization and/or infection
680 due to *Candida* spp [151].

681 More recent studies have examined this approach in high-risk patients,
682 particularly those undergoing hemodialysis [194-197]. Three randomized, controlled
683 trials have evaluated the use of 10% povidone iodine [195-197]. A significant decrease
684 in colonization, exit-site infection, or bloodstream infection was observed. The beneficial
685 effect was most prominent in subjects with nasal colonization by *S. aureus* [195-197].

686 Nasal carriers of *S. aureus* are more likely to experience a CRBSI than non-
687 colonized persons [201-203]. This has prompted investigators to assess the utility of
688 topical mupirocin, a potent anti-staphylococcal agent. Several studies have demonstrated
689 a reduced risk of CRBSI when mupirocin ointment was applied at the catheter insertion
690 site [195, 204-206]. Others have shown similar benefits when mupirocin was applied

691 [intran](#)asally [202, 203, 207]. However, enthusiasm for this measure has been dampened
by

692 the rapid emergence of mupirocin resistance observed at some centers [150, 208, 209],

693 and the potential degrading effect that mupirocin has on polyurethane catheters [154,

694 155].

695 In the only study demonstrating a significant effect on mortality, the application
696 of bacitracin/neomycin/polymyxin B ointment at the catheter insertion site was compared
697 to placebo in 169 hemodialysis patients [210]. Infections were observed in more patients
698 in the placebo group than in the bacitracin/neomycin/polymyxin B group (34 versus 12%;
699 relative risk, 0.35; 95% CI, 0.18 to 0.68; $P = 0.0013$). The number of infections per 1,000
700 catheter days (4.10 versus 1.02; $P < 0.0001$) and the number of bacteremias per 1,000
701 catheter days (2.48 versus 0.63; $P = 0.0004$) were also greater in the placebo group.

702 Within the 6-month study period, there were 13 deaths in the placebo group as compared
703 with three deaths in the bacitracin/neomycin/polymyxin B group ($P = 0.004$). Thus, there
704 is evidence from one study in hemodialysis patients that bacitracin/neomycin/polymyxin
705 B ointment can improve outcome, but no similar data exist for other patient populations
706 [210].

707 **Antibiotic Lock Prophylaxis, Antimicrobial Catheter Flush and Catheter Lock**

708 **Prophylaxis**

709 **Recommendation**

710 Use prophylactic antimicrobial lock solution in patients with long term catheters who
711 have a history of multiple CRBSI despite optimal maximal adherence to aseptic
712 technique [23, 211-228]. Category II

713 **Background**

714 To prevent CRBSI, a wide variety of antibiotic and antiseptic solutions have been
715 utilized to flush or lock catheter lumens [23, 211-228]. Catheter lock is a technique by
716 which an antimicrobial solution is used to fill a catheter lumen and then allowed to dwell
717 for a period of time while the catheter is idle. Antibiotics of various concentrations that
718 have been used either alone (when directed at a specific organism) or in combination (to
719 achieve broad empiric coverage) to prophylactically flush or lock central venous
720 catheters include vancomycin, gentamicin, ciprofloxacin, minocycline, amikacin,
721 cefazolin, cefotaxime, and ceftazidime; while antiseptics have included alcohol,
722 taurolidine, trisodium citrate. (Taurolidine and trisodium citrate are not approved for this
723 use in the US). These agents are usually combined with a compound acting as an
724 anticoagulant, such as heparin or ethylenediaminetetraacetic acid (EDTA). Most of these
725 studies have been conducted in relatively small numbers of high-risk patients, such as
726 hemodialysis patients, neonates, or neutropenic oncology patients. Although most
727 studies indicate a beneficial effect of the antimicrobial flush or lock solution in terms of
728 prevention of catheter-related infection, this must be balanced by the potential for side
729 effects, toxicity, allergic reactions, or emergence of resistance associated with the
730 antimicrobial agent. The wide variety of compounds used, the heterogeneity of the
731 patient populations studied, and limitations in the size or design of studies preclude a
732 general recommendation for use. In addition, there are no FDA-approved formulations
733 approved for marketing, and most formulations have been prepared in hospital
734 pharmacies. A brief overview of some of the studies follows.

735 At least 10 studies regarding catheter flush or lock solutions have been performed
736 in hemodialysis patients [218, 219, 221-228]. Three meta-analyses have all demonstrated
737 that catheter lock solutions reduce risk of CRBSI in hemodialysis patients [229-231]. In
738 the largest of these studies, 291 subjects were enrolled in a prospective randomized
739 comparison of 30% trisodium citrate versus heparin [223]. The rate of CRBSI was
740 significantly lower in the group whose catheters were locked with trisodium citrate (4.1
741 BSI/1,000 CVC days vs. 1.1 BSI/1,000 CVC days, $P < 0.001$), and no significant
742 difference in thrombosis or occlusion of the catheter was noted. However, if infused
743 rapidly, concentrated citrate can result in serious hypocalcaemia, cardiac dysrhythmia,
744 and death. The second largest study in hemodialysis subjects examined the effect of a
745 catheter lock solution containing cefazolin, gentamicin, and heparin compared to control
746 patients receiving only heparin [225]. In 120 subjects, the rate of CRBSI was
747 significantly lower in those receiving the antibiotic lock solution (0.44 BSI/1,000 CVC
748 days vs. 3.12 BSI/1,000 CVC days, $P = 0.03$) [225]. Other trials in hemodialysis patients
749 have studied minocycline, gentamicin, EDTA, heparin, taurolidine, vancomycin, and
750 cefotaxime.

751 At least five studies have been conducted in pediatric oncology patients [211, 212,
752 215-217]. In the largest trial, 126 subjects were enrolled in a prospective, randomized,
753 double blind study comparing vancomycin/ciprofloxacin/heparin (VCH) to
754 vancomycin/heparin (VH) to heparin (H) alone [215]. The time to CVC-related infection
755 was significantly longer in the VCH or VH arms of the study compared to heparin, and
756 the rate of possible or definite catheter-related infection was significantly lower with

757 either of the antibiotic containing solutions compared to heparin alone (1.72/1,000 CVC
758 days [H] vs. 0.55/1,000 CVC days [VCH] vs. 0.37/1,000 CVC days [VH]).

759 In a meta-analysis of seven randomized, controlled trials examining the utility of
760 vancomycin-containing lock or flush solutions compared to heparin alone, the risk ratio
761 for vancomycin/heparin solutions was 0.49 (95% CI 0.26-0.95, $p = 0.03$) [232]. Use of
762 the catheter lock technique appeared to have greater benefit than simply flushing
763 vancomycin through the catheter.

764 Recently, a prospective, double blind, randomized trial compared the utility of
765 70% ethanol lock versus heparinized saline for the prevention of CABSIs in oncology
766 patients. Patients receiving the ethanol lock preventive therapy were significantly less
767 likely to experience a CABSI (0.60/1,000 CVC days vs. 3.11/1,000 CVC days; OR 0.18,
768 95% CI 0.05-0.65, $p = 0.008$) [233].

769 Anticoagulants

770 Recommendation

771 Do not routinely use anticoagulant therapy to reduce the risk of catheter-related infection
772 in general patient populations [234]. Category II

773 Background

774 Shortly after insertion, intravascular catheters are coated with a conditioning film,
775 consisting of fibrin, plasma proteins, and cellular elements, such as platelets and red
776 blood cells [27, 235]. Microbes interact with the conditioning film to result in
777 colonization of the catheter [236]. There is a close association between thrombosis of
778 central venous catheters and infection [35, 237, 238]. Therefore, anticoagulants have
779 been used to prevent catheter thrombosis and presumably reduce the risk of infection.

780 In a meta-analysis evaluating the benefit of heparin prophylaxis (3 units/mL in
781 parenteral nutrition, 5,000 units every 6 or 12 hours flush or 2,500 units low molecular
782 weight heparin subcutaneously) in patients with short-term CVCs, the risk for catheter-
783 related central venous thrombosis was reduced with the use of prophylactic heparin
784 [234]. However, no substantial difference in the rate of CRBSI was observed. In a more
785 recent prospective, randomized trial, 204 patients with non-tunneled catheters were
786 assigned to receive a continuous infusion of heparin (100 units/kg/d) or saline (50 mL/d)
787 [239]. The rate of CRBSI was significantly decreased in the group receiving heparin (2.5
788 BSI/1,000 CVC days vs. 6.4 BSI/1,000 CVC days). Because the majority of heparin
789 solutions contain preservatives with antimicrobial activity, whether any decrease in the
790 rate of CRBSI is a result of the reduced thrombus formation, the preservative, or both is
791 unclear.

792 The majority of pulmonary artery, umbilical, and central venous catheters are
793 available as heparin-bonded devices. The majority of catheters are heparin bonded with
794 benzalkonium, which provides the catheters with antimicrobial activity [240] and
795 provides an anti-thrombotic effect [241]. However, some catheters have heparin bound
796 directly to the catheter without benzalkonium [242]. Studies have shown that heparin-
797 bonded catheters reduce risk of thrombosis and risk of CRBSI [239, 241-243]; but are
798 less effective at reducing catheter colonization than catheters impregnated with
799 chlorhexidine/silver sulfadiazine [244]. Unfortunately, heparin-induced
800 thrombocytopenia can occur and has prompted many clinicians to avoid heparin [245].
801 Trisodium citrate has been recommended as a catheter lock solution because it possesses
802 both anticoagulant and antimicrobial properties [223]. In a prospective, randomized,

803 double blind study in hemodialysis patients, use of interdialytic heparin (5,000 U/mL)
804 was associated with a significantly greater rate of CRBSIs compared to use of 30%
805 trisodium citrate (4.1 BSI/1,000 CVC days vs. 1.1BSI/1,000 CVC days [246].

806 Warfarin has been evaluated as a means to reduce CVC thrombus formation and,
807 hence, infection [247-251]. However, other studies have not confirmed reduced
808 thrombosis and others have found untoward interactions in patients receiving 5-FU [252,
809 253]. Data are quite limited; and although low dose warfarin decreases the risk of
810 thrombus formation in cancer patients, it has not been shown to reduce infectious
811 complications. Over 20% of patients in some studies develop prolonged prothrombin
812 times and required dosage adjustment [254]. Other anticoagulants, such as factor Xa
813 inhibitors or direct thrombin inhibitors, have not been adequately assessed in terms of
814 reducing the risk of catheter-associated infection.

815 **Replacement of Peripheral and Midline Catheters**

816 **Recommendations**

817 1. Replace peripheral catheters every 72-96 hours to reduce risk of infection and
818 phlebitis in adults. Category 1B
819 2. Replace peripheral catheters in children only when clinically indicated [82, 83].
820 Category 1B

821 2. Replace midline catheters only when there is a specific indication. Category II

822 **Background**

823 Scheduled replacement of intravascular catheters has been proposed as a method
824 to prevent phlebitis and catheter-related infections. Studies of short peripheral venous
825 catheters indicate that the incidence of thrombophlebitis and bacterial colonization of

826 catheters increases when catheters are left in place >72 hours [83, 255, 256]. However,
827 rates of phlebitis are not substantially different in peripheral catheters left in place 72
828 hours compared with 96 hours [257]. Because phlebitis and catheter colonization have
829 been associated with an increased risk for catheter-related infection, short peripheral
830 catheter sites commonly are replaced at 72-96 hour intervals to reduce both the risk for
831 infection and patient discomfort associated with phlebitis.

832 Midline catheters are associated with lower rates of phlebitis than short peripheral
833 catheters and with lower rates of infection than CVCs [258-260]. In one prospective
834 study of 140 midline catheters, their use was associated with a BSI rate of 0.8 per 1,000
835 catheter days [260]. No specific risk factors, including duration of catheterization, were
836 associated with infection. Midline catheters were in place a median of 7 days, but for as
837 long as 49 days. Although the findings of this study suggested that midline catheters
838 could be changed only when there is a specific indication, no prospective, randomized
839 studies have assessed the benefit of routine replacement as a strategy to prevent CRBSI
840 associated with midline catheters.

841 **Replacement of CVCs, Including PICCs and Hemodialysis Catheters**

842 **Recommendations**

843 1. Do not routinely replace CVCs, PICCs, hemodialysis catheters, or pulmonary artery
844 catheters to prevent catheter-related infections. Category IB

845 2. Do not remove CVCs or PICCs on the basis of fever alone. Use clinical judgment
846 regarding the appropriateness of removing the catheter if infection is evident elsewhere or
847 if a noninfectious cause of fever is suspected. Category II

Comment [jds21]: Suggest a statement about the importance of verifying that a CVL is truly a central line and not a midline re: fluids infused thru it and inclusion in denominator for CLABSI.

Comment [jds22]: Suggest: remove CVL if fungemia develops; for bacterial infections, remove if persistently bacteremic or hemodynamically unstable without other source identified.

848 3. Do not use guidewire exchanges routinely for non-tunneled catheters to prevent
849 infection. Category IB

850 4. Do not use guidewire exchanges to replace a non-tunneled catheter suspected of
851 infection. Category IB

852 4. Use a guidewire exchange to replace a malfunctioning non-tunneled catheter if no
853 evidence of infection is present. Category IB

854 5. Use new sterile gloves before handling the new catheter when guidewire exchanges are
855 performed. Category II

856 **Background**

857 Catheter replacement at scheduled time intervals as a method to reduce CRBSI
858 has not lowered rates. Two trials have assessed a strategy of changing the catheter every
859 7 days compared with a strategy of changing catheters as needed [261, 262]. One of these
860 studies involved 112 surgical ICU patients needing CVCs, pulmonary artery catheters, or
861 peripheral arterial catheters [262], whereas the other study involved only subclavian
862 hemodialysis catheters [261]. In both studies, no difference in CRBSI was observed in
863 patients undergoing scheduled catheter replacement every 7 days compared with patients
864 whose catheters were replaced as needed.

865 Scheduled guidewire exchanges of CVCs are another proposed strategy for
866 preventing CRBSI. The results of a meta-analysis of 12 randomized, controlled trials
867 assessing CVC management failed to prove any reduction of CRBSI rates through routine
868 replacement of CVCs by guidewire exchange compared with catheter replacement on an
869 as needed basis [263]. Thus, routine replacement of CVCs is not necessary for catheters
870 that are functioning and have no evidence of causing local or systemic complications.

871 Catheter replacement over a guidewire has become an accepted technique for
872 replacing a malfunctioning catheter or exchanging a pulmonary artery catheter for a CVC
873 when invasive monitoring no longer is needed. Catheter insertion over a guidewire is
874 associated with less discomfort and a significantly lower rate of mechanical
875 complications than are those percutaneously inserted at a new site [264]. In addition, this
876 technique provides a means of preserving limited venous access in some patients.

877 Replacement of temporary catheters over a guidewire in the presence of bacteremia is not
878 an acceptable replacement strategy because the source of infection is usually colonization
879 of the skin tract from the insertion site to the vein [25, 264]. However, in selected patients

880 with tunneled hemodialysis catheters and bacteremia, catheter exchange over a
881 guidewire, in combination with antibiotic therapy, is an alternative as a salvage strategy
882 in patients with limited venous access [265-268].

883 Because of the increased difficulty obtaining vascular access in children, attention
884 should be given to the frequency with which catheters are replaced in these patients. In a
885 study in which survival analysis techniques were used to examine the relation between
886 the duration of central venous catheterization and complications in pediatric ICU patients,
887 all of the patients studied ($n = 397$) remained uninfected for a median of 23.7 days [121].
888 In addition, no relation was found between duration of catheterization and the daily
889 probability of infection ($r = 0.21$; $p > 0.1$), suggesting that routine replacement of CVCs
890 likely does not reduce the incidence of catheter-related infection [121].

891 Vascular access sites can be even more limited among neonates. Four
892 randomized trials ($n=368$) summarized in a recent Cochrane Database Systemic Review
893 compared the effects of giving parenteral nutrition through percutaneous central venous

Formatted: Highlight

894 catheters vs. peripheral intravenous catheters. Fewer painful procedures (venopunctures)
895 were required in neonates randomized to percutaneously placed CVCs, and there was no
896 evidence for increased risk of BSIs [269].

897 CVC occlusion due to thrombus formation is one of the most common reasons for
898 CVC removal in neonates. Various methods have been tried to prevent catheter
899 occlusion. Recently, a randomized trial (n=201) evaluated whether a continuous heparin
900 infusion (0.5 units/kg/hour) could effectively prolong the duration of catheterization
901 when compared to a placebo infusion. The rate of catheter occlusion requiring catheter
902 removal was lower in the heparin group (6% vs. 31%, P=0.001; NNT=4). Rates of
903 CRBSI were similar, although the study was not powered to evaluate CRBSI rate
904 differences. Heparin associated antibody levels were not routinely measured [270].

905 **Hemodialysis Catheters**

906 The use of catheters for hemodialysis is the most common factor contributing to
907 bacteremia in dialysis patients [271, 272]. The relative risk for bacteremia in patients
908 with dialysis catheters is sevenfold the risk for patients with arteriovenous (AV) fistulas
909 [273]. To reduce the rate of infection, hemodialysis catheters should be avoided in favor
910 of AV fistulas and grafts. If temporary access is needed for dialysis, a cuffed catheter is
911 preferable to a non-cuffed catheter, even in the ICU setting, if the catheter is expected to
912 stay in place for >3 weeks [198].

913 **Pulmonary Artery Catheters**

914 Pulmonary artery catheters are inserted through a polytetrafluoroethylene
915 introducer and typically remain in place an average of 3 days. The majority of pulmonary
916 artery catheters are heparin bonded, which reduces not only catheter thrombosis but also

917 microbial adherence to the catheter [240]. Meta-analysis indicates that the CRBSI rate
918 associated with pulmonary artery catheterization is 3.7 per 1,000 catheter days and
919 somewhat higher than the rate observed for unmedicated and non-tunnelled CVCs (2.7
920 per 1,000 catheter days)[6, 93].

921 Data from prospective studies indicate that the risk of significant catheter
922 colonization and CRBSI increases the longer the catheter remains in place. In general, the
923 risk of significant catheter colonization increases after 4 days of catheterization [137,
924 274, 275], whereas the risk of CRBSI increases beyond 5-7 days of catheterization [137,
925 146, 276]. Efforts must be made to differentiate between infection related to the
926 introducer and that related to the pulmonary artery catheter. Significant colonization of
927 the introducer occurs earlier than that of the pulmonary artery catheter [274, 277].
928 However, no studies indicate that catheter replacement at scheduled time intervals is an
929 effective method to reduce CRBSI [262, 264, 277]. In patients who continue to require
930 hemodynamic monitoring, pulmonary artery catheters do not need to be changed more
931 frequently than every 7 days [277]. No specific recommendation can be made regarding
932 routine replacement of catheters that need to be in place for >7 days.

933 Pulmonary artery catheters are usually packaged with a thin plastic sleeve that
934 prevents touch contamination when placed over the catheter. In a study of 166 catheters,
935 patients who were randomly assigned to have their catheters self-contained within this
936 sleeve had a reduced risk for CRBSI compared with those who had a pulmonary artery
937 catheter placed without the sleeve ($p = 0.002$) [138].

938 **Umbilical Catheters**

939 **Recommendations**

- 940 1. Remove and do not replace umbilical artery catheters if any signs of CRBSI, vascular
941 insufficiency, or thrombosis are present [278]. Category II
- 942 2. Remove and do not replace umbilical venous catheters if any signs of CRBSI or
943 thrombosis are present [278]. Category II
- 944 3. No recommendation can be made for treating through an umbilical venous catheter
945 suspected of being infected. Unresolved issue
- 946 4. Replace umbilical venous catheters only if the catheter malfunctions. Category II
- 947 5. Cleanse the umbilical insertion site with an antiseptic before catheter insertion. Avoid
948 tincture of iodine because of the potential effect on the neonatal thyroid. Other iodine-
949 containing products (e.g., povidone iodine) can be used [279-283]. Category IB
- 950 6. Do not use topical antibiotic ointment or creams on umbilical catheter insertion sites
951 because of the potential to promote fungal infections and antimicrobial resistance [150,
952 151]. Category IA
- 953 7. Add low doses of heparin (0.25-1.0 U/ml) to the fluid infused through umbilical
954 arterial catheters [284-286]. Category IB
- 955 8. Remove umbilical catheters as soon as possible when no longer needed or when any
956 sign of vascular insufficiency to the lower extremities is observed. Optimally, umbilical
957 artery catheters should not be left in place >5 days [278, 287]. Category II
- 958 9. Umbilical venous catheters should be removed as soon as possible when no longer
959 needed, but can be used up to 14 days if managed aseptically [288, 289]. Category II

960 **Background**

961 Although the umbilical stump becomes heavily colonized soon after birth,
962 umbilical vessel catheterization often is used for vascular access in newborn infants.

963 Umbilical vessels can be cannulated easily and permit both collection of blood samples
964 and measurement of hemodynamic status. The incidences of catheter colonization and
965 BSI are similar for umbilical vein catheters and umbilical artery catheters. In several
966 studies, an estimated 40%-55% of umbilical artery catheters were colonized and 5%
967 resulted in CRBSI; umbilical vein catheters were associated with colonization in 22%-
968 59% of cases [280, 281, 290] and with CRBSI in 3%-8% of cases [281]. Although
969 CRBSI rates are similar for umbilical catheters in the high position (i.e., above the
970 diaphragm) compared with the low position (i.e., below the diaphragm and above the
971 aortic bifurcation), catheters placed in the high position result in a lower incidence of
972 vascular complications without an increase in adverse sequelae [281].

973 Risk factors for infection differ for umbilical artery and umbilical vein catheters.
974 In one study, neonates with very low birth weight who also received antibiotics for >10
975 days were at increased risk for umbilical artery CRBSIs [281]. In comparison, those with
976 higher birth weight and receipt of parenteral nutrition fluids were at increased risk for
977 umbilical vein CRBSI. Duration of catheterization was not an independent risk factor for
978 infection of either type of umbilical catheter.

979 A recent randomized trial (n=210) evaluated whether long-term umbilical venous
980 catheterization (up to 28 days) would result in the same or fewer CABSIs when compared
981 with neonates who were randomized to short-term umbilical venous catheterization for 7-10
982 days followed by percutaneous central venous catheterization. CABSIs rate was higher
983 (20%) among long term catheterized neonates when compared to short term catheterized
984 neonates (13%). The difference was not statistically significant ($P=0.17$), although the
985 study was underpowered to evaluate differences in venous thrombosis rates [291].

Deleted: to

986 Peripheral Arterial Catheters and Pressure Monitoring Devices for Adult and

987 Pediatric Patients

988 Recommendations

989 1. In adults, use of the radial, brachial or dorsalis pedis sites is preferred over the femoral
990 or axillary sites of insertion to reduce the risk of infection [94, 95, 292, 293]. Category IB

991 2. In children, the brachial site should not be used. The radial, dorsalis pedis, and
992 posterior tibial sites are preferred over the femoral or axillary sites of insertion [94].

993 Category II

994 3. A cap, mask, sterile gloves and a large sterile fenestrated drape should be used during

Formatted: Highlight

995 peripheral arterial catheter insertion [95, 293]. Category IB

996 4. During axillary or femoral artery catheter insertion, maximal sterile barriers
997 precautions should be used. Category II

998 5. Replace arterial catheters only when there is a clinical indication. Category II

999 6. Remove the arterial catheter as soon as it is no longer needed. Category II

1000 7. Use disposable, rather than reusable, transducer assemblies when possible [294-298].

1001 Category IB

1002 8. Do not routinely replace arterial catheters to prevent catheter-related infections [262,
1003 276, 299, 300]. Category II

1004 9. Replace disposable or reusable transducers at 96-hour intervals. Replace other
1005 components of the system (including the tubing, continuous-flush device, and flush

1006 solution) at the time the transducer is replaced [25, 295]. Category IB

1007 10. Keep all components of the pressure monitoring system (including calibration devices
1008 and flush solution) sterile [294, 301-303]. Category IA

1009 11. Minimize the number of manipulations of and entries into the pressure monitoring
1010 system. Use a closed flush system (i.e., continuous flush), rather than an open system
1011 (i.e., one that requires a syringe and stopcock), to maintain the patency of the pressure
1012 monitoring catheters [297, 304]. Category II

1013 12. When the pressure monitoring system is accessed through a diaphragm, rather than a
1014 stopcock, wipe the diaphragm with an appropriate antiseptic before accessing the system
1015 [297]. Category IA

1016 13. Do not administer dextrose-containing solutions or parenteral nutrition fluids through
1017 the pressure monitoring circuit [297, 305, 306]. Category IA

1018 14. Sterilize reusable transducers according to the manufacturers' instructions if the use of
1019 disposable transducers is not feasible [297, 305-308]. Category IA

1020 **Background**

1021 Peripheral arterial catheters are usually inserted into the radial or femoral artery
1022 and permit continuous blood pressure monitoring and blood gas measurements. The rate
1023 of CRBSI is lower than that of short term, uncuffed, non-coated, non-tunneled CVCs (1.7
1024 versus 2.7 per 1,000 catheter days)[6]. However, CRBSI rates are comparable between
1025 arterial catheters and short term, uncuffed, medicated, non-tunneled CVCs [6]. Unlike
1026 CVCs, use of full barrier precautions during arterial cannulation does not appear to reduce
1027 the risk of arterial CRBSI [293, 309]. Nonetheless, when arterial catheters are inserted
1028 using a protocol which includes maximum barrier precautions, a very low rate of CRBSI
1029 (0.41/1,000 catheter days) can be achieved[95]. Although a meta-analysis failed to
1030 discern a difference in rates of CRBSI among three sites of insertion (radial, femoral, and
1031 axillary)[310], colonization of catheters inserted in the femoral site occurs more often

Formatted: Highlight

1032 [293]. In addition, a prospective observational study of over 2,900 arterial catheters that
1033 were inserted using maximum barrier precautions demonstrated an almost 8-fold increase
1034 in the incidence of CRBSI when the femoral site was used compared to the radial
1035 site[311]. Furthermore, there is a greater risk of CRBSI caused by Gram-negative
1036 bacteria when the femoral site is utilized [311]. The rates of catheter colonization and
1037 CRBSI appear similar between the radial and dorsalis pedis sites[292]. The risk of
1038 developing a CRBSI increases with the duration of catheterization [276, 312]; however,
1039 the routine changing of arterial catheters at scheduled times does not result in a
1040 diminution of the rate of CRBSI [262]. Catheters that need to be in place for >5 days
1041 should not be routinely changed if no evidence of infection is observed.

1042 **Replacement of Administration Sets**

1043 **Recommendations**

- 1044 1. In patients not receiving blood, blood products or lipid emulsions, replace
1045 administration sets, including secondary sets and add-on devices, no more frequently than
1046 at 96-hour intervals, [313] but at least every 7 days [255, 314-316]. Category IA
- 1047 2. Replace tubing used to administer blood, blood products, or lipid emulsions (those
1048 combined with amino acids and glucose in a 3-in-1 admixture or infused separately)
1049 within 24 hours of initiating the infusion [317-320]. Category IB
- 1050 3. Replace tubing used to administer propofol infusions every 6 or 12 hours, when the
1051 vial is changed, per the manufacturer's recommendation (FDA website Medwatch) [321].
1052 Category IA

1053 **Background**

1054 The optimal interval for routine replacement of IV administration sets has been
1055 examined in a number of well-controlled studies and meta-analyses. Data from these
1056 studies reveal that replacing administration sets no more frequently than 72-96 hours after
1057 initiation of use is safe and cost-effective [255, 257, 313, 315, 316]. More recent studies
1058 suggest that administration sets may be used safely for up to 7 days if used in conjunction
1059 with antiseptic catheters or if fluids that enhance microbial growth (e.g., parenteral
1060 nutrition or blood) have not been used [30, 322]. When a fluid that enhances microbial
1061 growth is infused (e.g., lipid emulsions and blood products), more frequent changes of
1062 administration sets are indicated as these products have been identified as independent
1063 risk factors for CRBSI [30, 317, 323-327].

1064 **Needleless Intravascular Catheter Systems**

1065 **Recommendations**

1066 1. Change the needleless components at least as frequently as the administration set.
1067 There is no benefit to changing these more frequently than every 72 hours [87, 328-334].

1068 Category II

1069 2. Change caps no more frequently than every 72 hours for the purpose of reduced
1070 infection rates or according to manufacturers' recommendations[328, 330, 333, 334].

1071 Category II

1072 3. Ensure that all components of the system are compatible to minimize leaks and breaks
1073 in the system[335]. Category II

Comment [JDS23]: More discussion on cap changes is needed, e.g., before obtaining blood cultures, frequency with TPN with/without lipids, propofol, etc.

1074 4. Minimize contamination risk by wiping the access port with an appropriate antiseptic

1075 (chlorhexidine preferred) and accessing the port only with sterile devices [330, 333, 335].

Formatted: Highlight

1076 Category IA

1077 5. Use a needleless system to access IV tubing. Category IC

1078 6. When needleless systems are used, the split septum valve is preferred over the

Formatted: Highlight

1079 mechanical valve due to increased risk of infection [336-339]. Category II

1080 Background

1081 Stopcocks used for injection of medications, administration of IV infusions, and

1082 collection of blood samples represent a potential portal of entry for microorganisms into

1083 vascular access catheters and IV fluids. Stopcock contamination is common, occurring in

1084 45% and 50% in the majority of series. Whether such contamination is a substantial entry

Comment [JDS24]: Please add citations here.

1085 point of CRBSI has been difficult to prove. Nonetheless, stopcocks should be capped

Comment [JDS25]: Shouldn't this be included as a specific recommendation?

1086 when not being used.

1087 "Piggyback" systems are used as an alternative to stopcocks. However, they also

1088 pose a risk for contamination of the intravascular fluid if the device entering the rubber

1089 membrane of an injection port is exposed to air or comes into direct contact with

1090 nonsterile tape used to fix the needle to the port. Modified piggyback systems have the

1091 potential to prevent contamination at these sites [340].

1092 Attempts to reduce the incidence of sharp injuries and the resultant risk for

1093 transmission of bloodborne infections to healthcare personnel have led to the design and

1094 introduction of needleless infusion systems. There are several types of needleless

1095 connectors commercially available.

Deleted: on the market.

1096 The first type of needleless system connectors consisted of a split septum cap,
1097 which is accessed with a blunt cannula instead of a needle. Because of the large amount
1098 of space in the hub to accommodate the cannula, blood can easily backup into this space
1099 and occlude the catheter. A luer-activated device, which incorporates a valve preventing
1100 the outflow of fluid through the connector, was designed to eliminate this problem. Some

1101 luer devices require a cap to be attached to the valve when not in use, which can be
1102 difficult to maintain aseptically, and therefore they may be prone to contamination.

1103 Another type of second-generation needleless system addressed the occlusion issue by
1104 incorporating positive pressure or neutral displacement to either flush out aspirated blood
1105 or prevent its aspiration into infusion catheters. However, with the positive pressure the
1106 risk of occlusion may actually rise, as the valves are held open, allowing retrograde blood
1107 flow into the catheters.

1108 Many studies have shown that when the devices are used according to
1109 manufacturers' recommendations (i.e., appropriate disinfection prior to access), they do
1110 not substantially affect the incidence of CRBSI [328-335]. Use of “second-generation”
1111 needleless connectors or positive pressure mechanical valves, which reduce the backflow
1112 of blood after it is disengaged, appear to be effective in reducing hub colonization in
1113 some [341-343], but not all studies [344]. In one study [341], the incidence of CRBSI
1114 was reduced when the needleless connector was compared to standard stopcocks.
1115 Appropriate disinfectants must be used to prevent transmission of microbes through

1116 connectors [345]. Disinfection of the devices with chlorhexidine/alcohol solutions
1117 appears to be most effective in reducing colonization [342]. However, reports continue

1118 to be published of outbreaks of CRBSI, even when the second-generation connectors are
1119 used [336-339]. The physical and mechanical properties of second-generation connectors
1120 vary widely from device to device. Potential explanations for outbreaks associated with

1121 these devices include difficulty encountered in adequate disinfection of the surface of the
1122 connector due to physical characteristics of the plastic housing diaphragm interface, fluid
1123 flow properties (laminar vs. turbulent), internal surface area, potential fluid dead space,
1124 inadequate flushing of the device due to poor visualization of the fluid flow pathway in
1125 opaque devices, and the presence of internal corrugations that could harbor organisms,
1126 particularly if the catheters are used to access blood [338]. Additionally, a silver coated
1127 connector valve has been approved for marketing. However, there are no published
1128 randomized trials with this device and no recommendation can be made regarding its use.
1129 Likewise, an antiseptic-barrier cap has been studied in a laboratory setting and appears to
1130 be effective in preventing the entry of microorganisms [346], but has not yet been studied
1131 in a clinical trial.

1132 **Multidose Parenteral Medication Vials and Parenteral Fluids**

1133 **Recommendations**

- 1134 1. Mix all routine parenteral fluids in the pharmacy in a laminar flow hood using aseptic
1135 technique [347, 348]. Category IB
- 1136 2. Do not use any container of parenteral fluid that has visible turbidity, leaks, cracks,
1137 particulate matter, or if the manufacturer's expiration date has passed [348]. Category IB
- 1138 3. Use single dose vials for parenteral additives or medications when possible [348, 349].
1139 Category II

1140 4. Do not combine the leftover content of single use vials for later use [348, 349].
1141 Category IA

1142 5. If multidose vials are used, refrigerate multidose vials after they are opened if
1143 recommended by the manufacturer [348]. Category II

1144 6. Cleanse the access diaphragm of multidose vials with 70% alcohol before inserting a
1145 device into the vial [350]. Category IA

1146 7. Use a sterile device to access a multidose vial and avoid touch contamination of the
1147 device before penetrating the access diaphragm [351, 352]. Category IA

1148 8. Discard multidose vial if sterility is compromised [351, 352]. Category IA

1149 9. All multidose vials should be dated when 1st used and thereafter not used beyond the
1150 manufacturer's stated expiration period. Category IC

1151 10. Use the needle and syringe to access the multidose vial only once and to then discard both
1152 safely. This applies to each and every dose withdrawn from the vial [351, 352]. Category IA

1153 11. Complete the infusion of lipid-containing solutions (e.g., 3-in-1 solutions) within 24
1154 hours of hanging the solution [317, 318, 326, 327, 353]Category IB

1155 12. Complete the infusion of lipid emulsions alone within 12 hours of hanging the
1156 emulsion. If volume considerations require more time, the infusion should be completed
1157 within 24 hours [317, 326, 327]. Category IB

1158 13. Complete infusions of blood or other blood products within 4 hours of hanging the
1159 blood[354-357]. Category II

1160 14. No recommendation can be made for the hang time of other parenteral fluids.
1161 Unresolved issue

1162 **Background**

1163 Parenteral medications commonly are dispensed in multidose, parenteral
1164 medication vials that might be used for prolonged periods for one or more patients.
1165 Although the overall risk for extrinsic contamination of multidose vials is likely minimal
1166 [358], the consequences of contamination might result in life threatening infection [359-
1167 361]].Risk of contamination must be minimized by using one needle and one syringe one
1168 time only. Simply changing the needle and using the same syringe to access the vial is an
1169 unacceptable practice. Single use vials are intended for single use only (one puncture).
1170 They are frequently preservative free and pose a risk for contamination if they are

1171 punctured several times. This is particularly true with propofol, a drug that readily
1172 supports the growth of bacteria once contaminated.

1173 **Performance Improvement**

1174 **Recommendation**

1175 Use hospital-specific or collaborative-based performance improvement initiatives in
1176 which multifaceted strategies are "bundled" together improve compliance with evidence-
1177 based recommended practices [61, 108, 109, 362-366]. Category 1B

1178 **Background**

1179 Clinical decision makers, healthcare payers, and patient safety advocates
1180 emphasize the importance of translating research findings into everyday practice.
1181 Rigorous evaluations of CRBSI preventive practices using study designs with high
1182 internal validity and including study populations that optimize external validity remain
1183 necessary. Once practices have been determined to be effective and economically
1184 efficient, the next step is to implement these evidence-based practices so they become

1185 part of routine clinical care. Unfortunately, the use of evidence-based CRBSI preventive
1186 practices in U.S. hospitals remains suboptimal [367, 368]. In a national survey conducted
1187 in March 2005 of over 700 U.S. hospitals, approximately one quarter of U.S. hospitals
1188 indicated that either maximal sterile barrier precautions during central line insertion or
1189 chlorhexidine gluconate as site disinfectant, two practices widely recommended to
1190 prevent CRBSI, were not being used routinely [369]. Approximately 15% of U.S.
1191 hospitals reported routinely changing CVCs to prevent infection despite evidence that
1192 this practice should no longer be used [368, 369].

1193 Accordingly, investigators have attempted various approaches to better translate
1194 research findings and evidence-based recommendations into clinical practice. Numerous
1195 quality improvement studies have been published during the past several years that have
1196 used various methods, such as education of healthcare personnel, audit and feedback,
1197 organizational change, and clinical reminders [54-57, 108, 109, 363, 370-372]. The
1198 educational interventions, for example, primarily targeted hand hygiene, use of maximal
1199 sterile barriers during insertion, appropriate insertion site selection, proper site care using
1200 chlorhexidine gluconate, and prompt removal of unnecessary catheters. While a large
1201 number of before-and-after studies with a few using concurrent control groups [61, 109]
1202 have been published, no randomized, controlled trial evaluating a quality improvement
1203 strategy to prevent CRBSI has been reported [373]. The vast majority of before-and-after
1204 studies reported statistically significant decreases in CRBSI rates after a quality
1205 improvement strategy was implemented [373]. Additionally, both controlled trials also
1206 found statistically significant reductions of CRBSI in the intervention units compared to
1207 control units [61, 109].

1208 Investigators have also employed multifaceted approaches in which several
1209 strategies are bundled together to improve compliance with evidence-based guidelines
1210 [61, 108, 109]. One such collaborative cohort study [108] of 108 ICUs in Michigan
1211 targeted clinicians' use of five evidence-based practices: hand hygiene, maximum barrier
1212 precautions, chlorhexidine site disinfection, avoiding the femoral site, and removing
1213 unnecessary central venous catheters. In addition to educating clinicians about CRBSI
1214 prevention, interventions used included: 1) a central venous catheter cart that contained
1215 all the necessary supplies; 2) a checklist to ensure adherence to proper practices; 3)
1216 stoppage of procedures in non-emergent situations, if evidence-based practices were not
1217 being followed; 4) prompt removal of central catheters during daily patient rounds; 5)
1218 feedback to the clinical teams regarding the number of CRBSI episodes and overall rates;
1219 and 6) buy-in from the chief executive officers of the participating hospitals that
1220 chlorhexidine gluconate products/solutions would be stocked prior to study initiation.
1221 Using an interrupted time series design and multivariable regression, the investigators
1222 reported a statistically significant 66% decrease in CRBSI rates approximately 18 months
1223 after the intervention began [108]. Specific process and outcome measures for tracking
1224 and feedback (i.e. rate of central line infections, proportion of central lines placed with all
1225 or individual bundle elements performed AND documented) should be identified in
1226 individual institutions based on areas that have been identified for performance
1227 improvement.

Comment [JDS26]: This generally does not occur *during* rounds, but rather the decision is made during rounds.

1228 Finally, emphasis on the care and maintenance of catheters once they are in place
1229 should be a focus of performance improvement and quality assurance in all programs. A
1230 study to assess practice and staff knowledge of CVC post-insertion care and identify

1231 aspects of CVC care with potential for improvement revealed several areas of opportunity
1232 to improve post-insertion care [374]. Rates of breaches in catheter care and CRBSIs
1233 were calculated and statistical significance assumed when $P < 0.05$. Data were recorded
1234 from 151 CVCs in 106 patients giving a total of 721 catheter days. In all, 323 breaches in
1235 care were identified giving a failure rate of 44.8%, with significant differences between
1236 intensive care unit (ICU) and non-ICU wards ($P < 0.001$). Dressings (not intact) and caps
1237 and taps? (incorrectly placed) were identified as the major lapses in CVC care with 158
1238 and 156 breaches per 1000 catheter days, respectively. Interventions to improve
1239 reliability of care should focus on making the implementation of best practice easier to
1240 achieve.

Formatted: Highlight

1241

1242

General comments:

- 1) We are disappointed that this guideline does not address clinical situations that we currently struggle with (e.g., cap changes related to blood culture draws; care of implanted cvls (ports); ecmo catheters; intra-atrial catheters).
- 2) We would like to see a more comprehensive discussion of CLABSI risk factors that would include ostomies, trachs, or any potential contamination with body fluids.
- 3) No information provided related to line insertion carts, cap change kits, or dressing change kits as adjunctive measures to prevent CLABSIs.
- 4) Since the title implies that these guidelines would be applicable to the care of central lines outside of acute care hospitals, would like to see more guidance for standardization of catheter care by home health agencies.
- 5) Use active voice for all recommendations. Some are inconsistent.
- 6) Do not see the need to have the recommendations duplicated in the text.
- 7) Would like to see more discussion of the bundled practices, the daily goal sheet, auditing insertion and maintenance practices. Might even include auditing in interventional radiology and in the O.R. Sometimes the recs and the ratings are

more definitive than expected when reading the background discussion that presents conflicting data (e.g. for use of maximal sterile barrier precautions.)

Table 1. Catheters used for venous and arterial access.

Catheter Type	Entry Site	Length	Comments
Peripheral venous catheters	usually inserted in veins of forearm or hand	less than 3 inches	phlebitis with prolonged use; rarely associated with bloodstream infection
Peripheral arterial catheters	usually inserted in radial artery; can be placed in femoral, axillary, brachial, posterior tibial arteries	less than 3 inches	low infection risk
Midline catheters	inserted via the antecubital fossa into the proximal basilic or cephalic veins	3 to 8 inches	anaphylactoid reactions have been reported with catheters made of elastomeric hydrogel; does not enter central veins; lower rates of phlebitis than short peripheral catheters
Nontunneled central venous catheters	percutaneously inserted into central veins (subclavian, internal jugular, or femoral)	8 cm or longer depending on patient size	account for majority of CRBSI
Pulmonary artery	inserted through a Polytetrafluoroethylene usually heparin bonded; similar introducer in a central vein	30 cm or longer depending on patient size	rates of bloodstream infection
as risk	(subclavian, internal jugular, or femoral)		CVC; subclavian site preferred to reduce infection
Peripherally inserted central venous catheters (PICC)	inserted into basilic, cephalic or brachial veins and enter the superior vena cava	20 cm or longer depending on patient size	lower rate of infection nontunneled CVCs
Tunneled central venous catheters tract; infection than	implanted into subclavian, internal jugular or femoral veins	8 cm or longer depending on patient size	cuff inhibits migration of organisms into catheter size lower rate of nontunneled CVC
Totally implantable image;	tunneled beneath skin and have devices subcutaneous port accessed	8 cm or longer depending on patient size	lowest risk for CRBSI; improved patient self-
required for	with a needle; implanted in subclavian or internal jugular vein		no need for local catheter site care; surgery catheter removal
Umbilical catheters with	inserted into either umbilical	6 cm or less, depending risk for CRBSI similar	

1304		vein or umbilical artery	on patient size	catheters placed in
1305	umbilical			
1306		vein vs. artery		
1307				
1308				
1309				
1310				

DRAFT

1311
1312 TABLE 2. Pooled means and key percentile of the distribution of central-line
1313 associated bloodstream in infection rates among hospitals participating in the National
1314 Healthcare Safety Network, CDC, 2006 –2007. [15]
1315
1316

Type of Intensive care Unit	No. Units	No. CABSIs	Catheter-days	Pooled mean/ 1,000 catheter-days	Percentile				
					10%	25%	50%	75%	90%
Burn	22	239	42452	5.6	0	1.5	3.8	8.2	13.5
Coronary	121	373	181079	2.1	0	0	1.3	2.8	5.3
Surgical cardiothoracic	97	397	275194	1.4	0	0	1.2	1.9	3.4
Medical	144	1073	454839	2.4	0	0.6	1.9	3.6	5.3
Medical/surgical	104	692	342214	2.0	0	0.5	1.5	3.0	4.2
Major teaching		972	662489	1.5	0	0	0.6	2.0	3.6
Med/Surg All others	343								
Pediatric medical/surgical	71	404	140,848	2.9	0.0	0.0	2.1	3.8	6.0
Neurologic	15	31	25440	1.2	-	-	-	-	-
Neurosurgical	39	173	68550	2.5	0	0	1.9	3.8	6.2
Surgical	128	881	383126	2.3	0	0.5	1.7	3.1	5.1
Trauma	32	435	107620	4.0	0.3	1.5	4.0	5.7	7.7
Inpatient medical ward	40	111	60257	1.8	0	0	0	2.2	3.4
Inpatient medical/surgical ward	82	169	132133	1.3	0	0	0	1.6	4.0

1317
1318

1319
1320
1321
1322

Comment [JDS27]: There are 2
Table 3s.

Table 3. Pooled means and key percentiles for the distribution of central-line associated
bloodstream infection rates for level III NICUs, NHSH, CDC, 2006-2007.[15]

Birth-weight category	No. units	No. CLABSI	Central line- days	Pooled mean	Percentile				
					10%	25%	50%	75%	90%
< 750 g	82	225	60850	3.7	0.0	0.0	2.3	4.9	9.0
751-1000 g	84	185	55445	3.3	0.0	0.0	2.4	4.5	7.3
1001-1500 g	83	144	55874	2.6	0.0	0.0	1.6	3.6	6.1
1501-2500 g	71	105	44402	2.4	0.0	0.0	1.1	3.3	6.0
>2500 g	61	87	42611	2.0	0.0	0.0	0.0	3.1	5.4

1323
1324

DRAFT

1325 TABLE 3. Most common pathogens isolated from nosocomial bloodstream infections,
1326 SCOPE. [16, 17]
1327

Pathogen	Percentage of BSIs		
	Total	ICU	Non-ICU
Coagulase-negative staphylococci	31.3	35.9	26.6
<i>Staphylococcus aureus</i>	20.2	16.8	23.7
Enterococcus spp.	9.4	9.8	9.0
<i>Candida</i> spp.	9.0	10.1	7.9
Gram-negative rods			
<i>Escherichia coli</i>	5.6	3.7	7.6
<i>Klebsiella</i> spp	4.8	4.0	5.5
<i>Enterobacter</i> spp.	4.3	4.7	3.8
<i>Pseudomonas aeruginosa</i>	3.9	4.7	3.1
<i>Acinetobacter baumannii</i>	1.7	2.1	1.3
<i>Serratia</i> spp.	1.3	1.6	0.9

1328

1329

DRAFT

1330
1331
1332
1333
1334
1335

DRAFT

1336
1337
1338 **Appendix B. Disclosure of financial interests or relationships.**
1339

1340 Naomi P. O'Grady, M.D.: No disclosures

1341 Mary Alexander, R.N.: No Disclosures

1342 Lillian A. Burns, M.T., M.P.H., C.I.C.

1343 E. Patchen Dellinger, M.D.

1344 Jeffery Garland, M.D.

1345 Stephen O. Heard, M.D.: Merck advisory board; Angiotech advisory board: Honoraria

1346 from Lippencott Wilkins and Williams;

1347 Pamela A. Lipsett, M.D.: No disclosures

1348 Henry Masur, M.D.:

1349 Leonard A. Mermel, D.O., Sc.M.

1350 Michele L. Pearson, M.D.:

1351 Issam I. Raad, M.D.

1352 Adrienne Randolph, M.D., M.Sc.

1353 Mark E. Rupp, M.D.

1354 Sanjay Saint, M.D., M.P.H.

1355

1356 Appendix C. Summary Recommendations

1357 **Strategies for Prevention of Catheter-Related Infections in Adult and Pediatric**

1358 **Patients**

1359 **Education, training and staffing**

1360 **Recommendations**

- 1361 1. Educate healthcare personnel regarding the indications for intravascular catheter use,
1362 proper procedures for the insertion and maintenance of intravascular catheters, and
1363 appropriate infection control measures to prevent intravascular catheter-related infections
1364 [53-61]. Category IA
- 1365 2. Periodically assess knowledge of and adherence to guidelines for all persons who are
1366 involved in the insertion and maintenance of intravascular catheters [53-61]. Category IA
- 1367 3. Designate only trained personnel who demonstrate competence for the insertion and
1368 maintenance of peripheral and central intravascular catheters. [60-74]. Category IA
- 1369 4. Ensure appropriate nursing staff levels in ICUs to minimize the incidence of catheter-
1370 related BSIs. Observational studies suggest a ratio of 2:1 in ICUs where nurses are
1371 managing patients with CVCs [75-77]. Category IB

1372 **Site selection**

1373 **Recommendations for peripheral catheters and midline catheters**

- 1374 1. In adults, use an upper-extremity site for catheter insertion. Replace a catheter inserted
1375 in a lower extremity site to an upper extremity site as soon as possible [82, 83]. Category
1376 IB
- 1377 2. In pediatric patients, the upper or lower extremities or the scalp can be used as the
1378 catheter insertion site [82, 83]. Category II

1379 3. Select catheters on the basis of the intended purpose and duration of use, known
1380 infectious and non-infectious complications (e.g., phlebitis and infiltration), and
1381 experience of individual catheter operators [83-85]. Category IB

1382 4. Avoid the use of steel needles for the administration of fluids and medication that
1383 might cause tissue necrosis, if extravasation occurs [83-85]. Category IA

1384 5. Use a midline catheter or peripherally inserted central catheter (PICC), instead of a
1385 short peripheral catheter, when the duration of IV therapy will likely exceed six days [83-
1386 85]. Category IB

1387 **Recommendations for central venous catheters**

1388 6. Weigh the risk and benefits of placing a central venous device at a recommended site
1389 to reduce infectious complications against the risk for mechanical complications (e.g.,
1390 pneumothorax, subclavian artery puncture, subclavian vein laceration, subclavian vein
1391 stenosis, hemothorax, thrombosis, air embolism, and catheter misplacement) [25, 86-
1392 101]. Category IA

1393 7. Use a subclavian site, rather than a jugular or a femoral site, in adult patients to
1394 minimize infection risk for nontunneled CVC placement [25, 99, 100]. Category IA

1395 8. No recommendation can be made for a preferred site of insertion to minimize infection
1396 risk for a tunneled CVC. Unresolved issue

1397 9. Place catheters used for hemodialysis and pheresis in a jugular or femoral vein, rather
1398 than a subclavian vein, to avoid venous stenosis [101-105]. Category IA

1399 10. Use ultrasound guidance to place central venous catheters to reduce the number of
1400 cannulation attempts and mechanical complications if this technology is available [106,
1401 107]. Category IB

1402 11. Promptly remove any intravascular catheter that is no longer essential [108, 109].

1403 Category IA

1404 **Hand Hygiene and Aseptic Technique**

1405 **Recommendations**

1406 1. Perform hand hygiene procedures, either by washing hands with conventional

1407 antiseptic containing soap and water or with waterless alcohol-based hand rubs (ABHR).

1408 Hand hygiene should be performed before and after palpating catheter insertion sites as

1409 well as before and after inserting, replacing, accessing, repairing, or dressing an

1410 intravascular catheter. Palpation of the insertion site should not be performed after the

1411 application of antiseptic, unless aseptic technique is maintained [58, 127-131]. Category

1412 IA

1413 2. Maintain aseptic technique for the insertion and care of intravascular catheters [25,

1414 132-134]. Category IA

1415 3. Wear clean gloves, rather than sterile gloves, for the insertion of peripheral

1416 intravascular catheters, if the access site is not touched after the application of skin

1417 antiseptics. Category IC

1418 4. Wear sterile gloves for the insertion of arterial, central, and midline

1419 catheters [25, 132-134]; and these gloves should be changed, if a catheter is being

1420 exchanged over a guidewire (thereby contaminating the gloves) and a new sterile catheter

1421 is then handled. Category IA

Deleted: S

Deleted: should be worn

1422 4. Wear either clean or sterile gloves when changing the dressing on intravascular
1423 catheters. Category IC

1424 **Maximal Sterile Barrier Precautions**

1425 **Recommendations**

1426 1. Use maximal sterile barrier precautions, including the use of a cap, mask, sterile gown,
1427 sterile gloves, and a large sterile full body drape, for the insertion of CVCs, PICCs, or
1428 guidewire exchange [60, 132, 136, 137]. Category IB

Comment [JDS28]: ? and peripheral
arterial catheters

1429 2. Use a sterile sleeve to protect pulmonary artery catheters during insertion [138].

1430 Category IB

1431 **Skin Preparation**

1432 **Recommendations**

1433

1434 1. Prepare clean skin with 70% alcohol before peripheral venous catheter insertion [139].

1435 Category IA

1436 2. Prepare clean skin site with a 2% chlorhexidine-based preparation before central
1437 venous catheter insertion and during dressing changes. If there is a contraindication to
1438 chlorhexidine, tincture of iodine, an iodophor, or 70% alcohol can be used as alternatives
1439 [140, 141]. Category IA

1440 3. No recommendation can be made for the safety or efficacy of chlorhexidine in infants
1441 aged <2 months. Unresolved issue

1442 4. Allow povidone iodine to remain on the skin for at least 2 minutes or longer for the
1443 antibacterial properties to take effect, if it is not yet dry before catheter insertion. The
1444 antibacterial properties of chlorhexidine work on contact, and chlorhexidine does not

1445 require a minimum 2- minute drying time before proceeding. Catheter insertion may
1446 begin as soon as the chlorhexidine is dry[140, 141]. Category IB

1447 **Catheter site dressing regimens**

1448 **Recommendations**

- 1449 1. Use either sterile gauze or sterile, transparent, semi-permeable dressing to cover the
1450 catheter site [146-149]. Category IA
- 1451 2. If the patient is diaphoretic or if the site is bleeding or oozing, use gauze dressing until
1452 this is resolved [146-149]. Category II
- 1453 3. Replace catheter site dressing if the dressing becomes damp, loosened, or visibly soiled
1454 [146, 147]. Category IB
- 1455 4. Do not use topical antibiotic ointment or creams on insertion sites, except for dialysis
1456 catheters, because of their potential to promote fungal infections and antimicrobial
1457 resistance [150, 151]. Category IB
- 1458 5. Do not submerge the catheter or catheter site in water. Showering should be permitted
1459 if precautions can be taken to reduce the likelihood of introducing organisms into the
1460 catheter (e.g., if the catheter and connecting device are protected with an impermeable
1461 cover during the shower) [152, 153]. Category II
- 1462 6. Replace dressings used on short-term CVC sites every 2 days for gauze dressings and
1463 at least every 7 days for transparent dressings, except in those pediatric patients in which
1464 the risk for dislodging the catheter may outweigh the benefit of changing the dressing
1465 [149]. Category IB
- 1466 7. Replace dressings used on tunneled or implanted CVC sites no more than once per
1467 week, until the insertion site has healed [149]Category IB

1468 8. No recommendation can be made regarding the necessity for any dressing on well-
1469 healed exit sites of long-term cuffed and tunneled CVCs. Unresolved issue

1470 9. Ensure that catheter site care is compatible with the catheter material [154, 155].

1471 Category IB

1472 10. Use a sterile sleeve for all pulmonary artery catheters [138]. Category IB

1473 11. Use a chlorhexidine-impregnated sponge dressing for temporary short-term catheters

1474 in patients older than 2 months of age, if the CRBSI rate is higher than the institutional

1475 goal, despite adherence to basic CRBSI prevention measures, including education and

1476 training, use of chlorhexidine for skin antisepsis, and MSB [22, 156-158]. Category 1B

1477 **Patient Cleansing**

1478 **Recommendation**

1479 Use a 2% chlorhexidine wash daily to reduce CRBSI [162]. Category II

1480 **Catheter Securement Devices**

1481 **Recommendation**

1482 Use a sutureless securement device to reduce the risk of infection for PICCs [163].

1483 Category II

1484 **Antimicrobial/Antiseptic Impregnated Catheters and Cuffs**

1485 **Recommendation**

1486 Use a chlorhexidine/silver sulfadiazine or minocycline/rifampin -impregnated CVC in

1487 adults whose catheter is expected to remain in place >5 days if, after successful

1488 implementation of a comprehensive strategy to reduce rates of CRBSI, the CRBSI rate

1489 remains above the goal set by the individual institution based on benchmark rates (Tables

1490 2 and 3) and local factors. The comprehensive strategy should include at least the

1491 following three components: educating persons who insert and maintain catheters, use of
1492 maximal sterile barrier precautions, and a 2% chlorhexidine preparation for skin
1493 antisepsis during CVC insertion. Category IA

1494 **Systemic Antibiotic Prophylaxis**

1495 **Recommendation**

1496 Do not administer systemic antimicrobial prophylaxis routinely before insertion or during
1497 use of an intravascular catheter to prevent catheter colonization or CRBSI [188].

1498 Category IA

1499 **Antibiotic Lock Prophylaxis, Antimicrobial Catheter Flush and Catheter Lock** 1500 **Prophylaxis**

1501 **Recommendation**

1502 Use prophylactic antimicrobial lock solution in patients with long term catheters who
1503 have a history of multiple CRBSI despite optimal maximal adherence to aseptic
1504 technique [23, 211-228]. Category II

1505 **Anticoagulants**

1506 **Recommendation**

1507 Do not routinely use anticoagulant therapy to reduce the risk of catheter-related infection
1508 in general patient populations [234]. Category II

1509 **Replacement of Peripheral and Midline Catheters**

1510 **Recommendations**

1511 1. Replace peripheral catheters every 72-96 hours to reduce risk of infection and
1512 phlebitis in adults. Category 1B

1513 2. Replace peripheral catheters in children only when clinically indicated [82, 83].

1514 Category 1B

1515 2. Replace midline catheters only when there is a specific indication. Category II

1516 **Replacement of CVCs, Including PICCs and Hemodialysis Catheters**

1517 **Recommendations**

1518 1. Do not routinely replace CVCs, PICCs, hemodialysis catheters, or pulmonary artery

1519 catheters to prevent catheter-related infections. Category IB

1520 2. Do not remove CVCs or PICCs on the basis of fever alone. Use clinical judgment

1521 regarding the appropriateness of removing the catheter if infection is evidenced

1522 elsewhere or if a noninfectious cause of fever is suspected. Category II

1523 3. Do not use guidewire exchanges routinely for non-tunneled catheters to prevent

1524 infection. Category IB

1525 4. Do not use guidewire exchanges to replace a non-tunneled catheter suspected of

1526 infection. Category IB

1527 4. Use a guidewire exchange to replace a malfunctioning non-tunneled catheter if no

1528 evidence of infection is present. Category IB

1529 5. Use new sterile gloves before handling the new catheter when guidewire exchanges are

1530 performed. Category II

1531 **Umbilical Catheters**

1532 **Recommendations**

1533 1. Remove and do not replace umbilical artery catheters if any signs of CRBSI, vascular

1534 insufficiency, or thrombosis are present [278]. Category II

1535 2. Remove and do not replace umbilical venous catheters if any signs of CRBSI or
1536 thrombosis are present [278]. Category II

1537 3. No recommendation can be made for treating through an umbilical venous catheter
1538 suspected of being infected. Unresolved issue

1539 4. Replace umbilical venous catheters only if the catheter malfunctions. Category II

1540 5. Cleanse the umbilical insertion site with an antiseptic before catheter insertion. Avoid
1541 tincture of iodine because of the potential effect on the neonatal thyroid. Other iodine-
1542 containing products (e.g., povidone iodine) can be used [279-283]. Category IB

1543 6. Do not use topical antibiotic ointment or creams on umbilical catheter insertion sites
1544 because of the potential to promote fungal infections and antimicrobial resistance [150,
1545 151]. Category IA

1546 7. Add low doses of heparin (0.25-1.0 U/ml) to the fluid infused through umbilical
1547 arterial catheters [284-286]. Category IB

1548 8. Remove umbilical catheters as soon as possible when no longer needed or when any
1549 sign of vascular insufficiency to the lower extremities is observed. Optimally, umbilical
1550 artery catheters should not be left in place >5 days [278, 287]. Category II

1551 9. Umbilical venous catheters should be removed as soon as possible when no longer
1552 needed, but can be used up to 14 days if managed aseptically [288, 289]. Category II

1553 **Peripheral Arterial Catheters and Pressure Monitoring Devices for Adult and**
1554 **Pediatric Patients**

1555 **Recommendations**

1556 1. In adults, use of the radial, brachial or dorsalis pedis sites is preferred over the femoral
1557 or axillary sites of insertion to reduce the risk of infection [94, 95, 292, 293]. Category IB

1558 2. In children, do not use the brachial site. The radial, dorsalis pedis, and
1559 posterior tibial sites are preferred over the femoral or axillary sites of insertion [94].

Deleted: should not be used

1560 Category II

1561 3. Use a cap, mask, sterile gloves and a large sterile fenestrated drape during
1562 peripheral arterial catheter insertion [95, 293]. Category IB

Deleted: A

Deleted: should be used

1563 4. Use maximal sterile barriers precautions during axillary or femoral artery catheter
1564 insertion. Category II

Deleted: .

Deleted: D

Deleted: . maximal sterile barriers||
1564 precautions should be used.

1565 5. Replace arterial catheters only when there is a clinical indication. Category II

1566 6. Remove the arterial catheter as soon as it is no longer needed. Category II

1567 7. Use disposable, rather than reusable, transducer assemblies when possible [294-298].

1568 Category IB

1569 8. Do not routinely replace arterial catheters to prevent catheter-related infections [262,
1570 276, 299, 300]. Category II

1571 9. Replace disposable or reusable transducers at 96-hour intervals. Replace other

1572 components of the system (including the tubing, continuous-flush device, and flush

1573 solution) at the time the transducer is replaced [25, 295]. Category IB

1574 10. Keep all components of the pressure monitoring system (including calibration devices

1575 and flush solution) sterile [294, 301-303]. Category IA

1576 11. Minimize the number of manipulations of and entries into the pressure monitoring

1577 system. Use a closed flush system (i.e., continuous flush), rather than an open system

1578 (i.e., one that requires a syringe and stopcock), to maintain the patency of the pressure

1579 monitoring catheters [297, 304]. Category II

1580 12. When the pressure monitoring system is accessed through a diaphragm, rather than a
1581 stopcock, wipe the diaphragm with an appropriate antiseptic before accessing the system
1582 [297]. Category IA

1583 13. Do not administer dextrose-containing solutions or parenteral nutrition fluids through
1584 the pressure monitoring circuit [297, 305, 306]. Category IA

1585 14. Sterilize reusable transducers according to the manufacturers' instructions if the use of
1586 disposable transducers is not feasible [297, 305-308]. Category IA

1587 **Replacement of Administration Sets**

1588 **Recommendations**

1589 1. In patients not receiving blood, blood products or lipid emulsions, replace
1590 administration sets, including secondary sets and add-on devices, no more frequently than
1591 at 96-hour intervals, [313] but at least every 7 days [255, 314-316]. Category IA

1592 2. Replace tubing used to administer blood, blood products, or lipid emulsions (those
1593 combined with amino acids and glucose in a 3-in-1 admixture or infused separately)
1594 within 24 hours of initiating the infusion [317-320]. Category IB

1595 3. Replace tubing used to administer propofol infusions every 6 or 12 hours, when the
1596 vial is changed, per the manufacturer's recommendation (FDA website Medwatch) [321].
1597 Category IA

1598 **Needleless Intravascular Catheter Systems**

1599 **Recommendations**

1600 1. Change the needleless components at least as frequently as the administration set.
1601 There is no benefit to changing these more frequently than every 72 hours [87, 328-334].
1602 Category II

1603 2. Change caps no more frequently than every 72 hours for the purpose of reduced
1604 infection rates or according to manufacturers' recommendations[328, 330, 333, 334].
1605 Category II

1606 3. Ensure that all components of the system are compatible to minimize leaks and breaks
1607 in the system[335]. Category II

1608 4. Minimize contamination risk by wiping the access port with an appropriate antiseptic
1609 (chlorhexidine preferred) and accessing the port only with sterile devices [330, 333, 335].
1610 Category IA

1611 5. Use a needleless system to access IV tubing. Category IC

1612 6. When needleless systems are used, the split septum valve is preferred over the
1613 mechanical valve due to increased risk of infection [336-339]. Category II

1614 **Multidose Parenteral Medication Vials and Parenteral Fluids**

1615 **Recommendations**

1616 1. Mix all routine parenteral fluids in the pharmacy in a laminar flow hood using aseptic
1617 technique [347, 348]. Category IB

1618 2. Do not use any container of parenteral fluid that has visible turbidity, leaks, cracks,
1619 particulate matter, or if the manufacturer's expiration date has passed [348]. Category IB

1620 3. Use single dose vials for parenteral additives or medications when possible [348, 349].
1621 Category II

1622 4. Do not combine the leftover content of single use vials for later use [348, 349].
1623 Category IA

1624 5. If multidose vials are used, refrigerate multidose vials after they are opened if
1625 recommended by the manufacturer [348]. Category II

1626 6. Cleanse the access diaphragm of multidose vials with 70% alcohol before inserting a
1627 device into the vial [350]. Category IA

1628 7. Use a sterile device to access a multidose vial and avoid touch contamination of the
1629 device before penetrating the access diaphragm [351, 352]. Category IA

1630 8. Discard multidose vial if sterility is compromised [351, 352]. Category IA

1631 9. All multidose vials should be dated when 1st used and thereafter not used beyond the
1632 manufacturer's stated expiration period. Category IC

1633 10. Use the needle and syringe to access the multidose vial only once and to then discard both
1634 safely. This applies to each and every dose withdrawn from the vial [351, 352]. Category IA

1635 11. Complete the infusion of lipid-containing solutions (e.g., 3-in-1 solutions) within 24
1636 hours of hanging the solution [317, 318, 326, 327, 353]Category IB

1637 12. Complete the infusion of lipid emulsions alone within 12 hours of hanging the
1638 emulsion. If volume considerations require more time, the infusion should be completed
1639 within 24 hours [317, 326, 327]. Category IB

1640 13. Complete infusions of blood or other blood products within 4 hours of hanging the
1641 blood[354-357]. Category II

1642 14. No recommendation can be made for the hang time of other parenteral fluids.
1643 Unresolved issue

1644 **Performance Improvement**

1645 **Recommendation**

1646 Use hospital-specific or collaborative-based performance improvement initiatives in
1647 which multifaceted strategies are "bundled" together improve compliance with evidence-
1648 based recommended practices [61, 108, 109, 362-366]. Category 1B

1649

1650

1651

1652

1653

1654

1655

1656

1657

1658

1659

1660

1661

1662

1663

1664

1665

1666

DRAFT

1667 References

1668

1669 1. Mermel LA. Prevention of intravascular catheter-related infections (Erratum: Ann

1670 Intern Med 133:395, 2000). Ann Intern Med 2000;132:391-402

Formatted: Swedish (Sweden)

1671 2. Dimick JB, Pelz RK, Consunji R, Swoboda SM, Hendrix CW and Lipsett PA.

1672 Increased resource use associated with catheter-related bloodstream infection in the

1673 surgical intensive care unit. Arch Surg 2001;136:229-34.

1674 3. Warren DK, Quadir WW, Hollenbeak CS, Elward AM, Cox MJ and Fraser VJ.

1675 Attributable cost of catheter-associated bloodstream infections among intensive care

1676 patients in a nonteaching hospital. Crit Care Med 2006;34:2084-9

Formatted: French (France)

1677 4. Blot SI, Depuydt P, Annemans L, et al. Clinical and economic outcomes in critically ill

1678 patients with nosocomial catheter-related bloodstream infections. Clin Infect Dis

1679 2005;41:1591-8

Formatted: French (France)

1680 5. Renaud B, Brun-Buisson C. Outcomes of primary and catheter-related bacteremia. A

1681 cohort and case-control study in critically ill patients. Am J Respir Crit Care Med

1682 2001;163:1584-90

1683 6. Maki DG, Kluger DM and Crnich CJ. The risk of bloodstream infection in adults with

1684 different intravascular devices: a systematic review of 200 published prospective studies.

1685 Mayo Clin Proc 2006;81:1159-71

Formatted: French (France)

1686 7. Bellini C, Petignat C, Francioli P, et al. Comparison of automated strategies for

1687 surveillance of nosocomial bacteremia. Infect Control Hosp Epidemiol 2007;28:1030-5

1688 8. Borlawsky T, Hota B, Lin MY, et al. Development of a reference information model
1689 and knowledgebase for electronic bloodstream infection detection. AMIA Annu Symp
1690 Proc 2008:56-60

Formatted: French (France)

1691 9. Rubin MA, Mayer J, Greene T, et al. An agent-based model for evaluating surveillance
1692 methods for catheter-related bloodstream infection. AMIA Annu Symp Proc 2008:631-5
1693 10. Trick WE, Zagorski BM, Tokars JI, et al. Computer algorithms to detect bloodstream
1694 infections. Emerg Infect Dis 2004;10:1612-20

1695 11. Horan TC, Andrus M and Dudeck MA. CDC/NHSN surveillance definition of health
1696 care-associated infection and criteria for specific types of infections in the acute care
1697 setting. Am J Infect Control 2008;36:309-32

Formatted: French (France)

1698 12. Mermel LA, Allon M, Bouza E, et al. Clinical practice guidelines for the diagnosis
1699 and management of intravascular catheter-related infection: 2009 Update by the
1700 Infectious Diseases Society of America. Clin Infect Dis 2009;49:1-45
1701 13. Joint Commission on the Accreditation of Healthcare Organizations. Accreditation
1702 manual for hospitals. In: Joint Commission on the Accreditation of Healthcare
1703 Organizations, ed. Chicago, IL, 1994:121-140

1704 14. National Nosocomial Infections Surveillance (NNIS) System Report, data summary
1705 from January 1992 through June 2004, issued October 2004. Am J Infect Control
1706 2004;32:470-85

1707 15. Edwards JR, Peterson KD, Andrus ML, Dudeck MA, Pollock DA and Horan TC.
1708 National Healthcare Safety Network (NHSN) Report, data summary for 2006 through
1709 2007, issued November 2008. Am J Infect Control 2008;36:609-26

1710 16. Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP and Edmond MB.
 1711 Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a
 1712 prospective nationwide surveillance study. Clin Infect Dis 2004;39:309-17

1713 17. Gaynes R, Edwards JR. Overview of nosocomial infections caused by gram-negative
 1714 bacilli. Clin Infect Dis 2005;41:848-54

1715 18. Burton DC, Edwards JR, Horan TC, Jernigan JA and Fridkin SK. Methicillin-
 1716 resistant Staphylococcus aureus central line-associated bloodstream infections in US
 1717 intensive care units, 1997-2007. JAMA 2009;301:727-36

1718 19. National Nosocomial Infections Surveillance (NNIS) system report, data summary
 1719 from January 1992-April 2000, issued June 2000. Am J Infect Control 2000;28:429-48

1720 20. Richards MJ, Edwards JR, Culver DH and Gaynes RP. Nosocomial infections in
 1721 medical intensive care units in the United States. National Nosocomial Infections
 1722 Surveillance System. Crit Care Med 1999;27:887-92

1723 21. Richards MJ, Edwards JR, Culver DH and Gaynes RP. Nosocomial infections in
 1724 pediatric intensive care units in the United States. National Nosocomial Infections
 1725 Surveillance System. Pediatrics 1999;103:103-9

1726 22. Garland JS, Alex CP, Mueller CD, et al. A randomized trial comparing povidone-
 1727 iodine to a chlorhexidine gluconate-impregnated dressing for prevention of central
 1728 venous catheter infections in neonates. Pediatrics 2001;107:1431-6.

1729 23. Garland JS, Alex CP, Henrickson KJ, McAuliffe TL and Maki DG. A vancomycin-
 1730 heparin lock solution for prevention of nosocomial bloodstream infection in critically ill
 1731 neonates with peripherally inserted central venous catheters: a prospective, randomized
 1732 trial. Pediatrics 2005;116:e198-205

Formatted: French (France)

Formatted: French (France)

Formatted: Portuguese (Brazil)

1733 24. Safdar N, Maki DG. The pathogenesis of catheter-related bloodstream infection with
 1734 noncuffed short-term central venous catheters. *Intensive Care Med* 2004;30:62-7

1735 25. Mermel LA, McCormick RD, Springman SR and Maki DG. The pathogenesis and
 1736 epidemiology of catheter-related infection with pulmonary artery Swan-Ganz catheters: a
 1737 prospective study utilizing molecular subtyping. *Am J Med* 1991;91:197S-205S

1738 26. Maki DG, Weise CE and Sarafin HW. A semiquantitative culture method for
 1739 identifying intravenous-catheter-related infection. *N Engl J Med* 1977;296:1305-9

1740 27. Raad I, Costerton W, Sabharwal U, Sacilowski M, Anaissie E and Bodey GP.
 1741 Ultrastructural analysis of indwelling vascular catheters: a quantitative relationship
 1742 between luminal colonization and duration of placement. *J Infect Dis* 1993;168:400-7

1743 28. Dobbins BM, Kite P, Kindon A, McMahon MJ and Wilcox MH. DNA fingerprinting
 1744 analysis of coagulase negative staphylococci implicated in catheter-related bloodstream
 1745 infections. *J Clin Pathol* 2002;55:824-8

1746 29. Anaissie E, Samonis G, Kontoyiannis D, et al. Role of catheter colonization and
 1747 infrequent hematogenous seeding in catheter-related infections. *Eur J Clin Microbiol*
 1748 *Infect Dis* 1995;14:134-7

1749 30. Raad I, Hanna HA, Awad A, et al. Optimal frequency of changing intravenous
 1750 administration sets: is it safe to prolong use beyond 72 hours? *Infect Control Hosp*
 1751 *Epidemiol* 2001;22:136-9.

1752 31. Mehall JR, Saltzman DA, Jackson RJ and Smith SD. Fibrin sheath enhances central
 1753 venous catheter infection. *Crit Care Med* 2002;30:908-12

1754 32. Donlan RM, Costerton JW. Biofilms: survival mechanisms of clinically relevant
 1755 microorganisms. *Clin Microbiol Rev* 2002;15:167-93

Formatted: French (France)

Formatted: Italian (Italy)

1756 33. Hawser SP, Douglas LJ. Biofilm formation by *Candida* species on the surface of
1757 catheter materials in vitro. *Infect Immun* 1994;62:915-21

1758 34. Stillman RM, Soliman F, Garcia L and Sawyer PN. Etiology of catheter-associated
1759 sepsis. Correlation with thrombogenicity. *Arch Surg* 1977;112:1497-9

1760 35. Raad, II, Luna M, Khalil SA, Costerton JW, Lam C and Bodey GP. The relationship
1761 between the thrombotic and infectious complications of central venous catheters. *JAMA*
1762 1994;271:1014-6

1763 36. Herrmann M, Suchard SJ, Boxer LA, Waldvogel FA and Lew PD. Thrombospondin
1764 binds to *Staphylococcus aureus* and promotes staphylococcal adherence to surfaces.
1765 *Infect Immun* 1991;59:279-88

1766 37. Shanks RM, Sargent JL, Martinez RM, Graber ML and O'Toole GA. Catheter lock
1767 solutions influence staphylococcal biofilm formation on abiotic surfaces. *Nephrol Dial*
1768 *Transplant* 2006;21:2247-55

1769 38. Chatzinikolaou I, Zipf TF, Hanna H, et al. Minocycline-ethylenediaminetetraacetate
1770 lock solution for the prevention of implantable port infections in children with cancer.
1771 *Clin Infect Dis* 2003;36:116-9

1772 39. McDevitt D, Francois P, Vaudaux P and Foster TJ. Identification of the ligand-
1773 binding domain of the surface-located fibrinogen receptor (clumping factor) of
1774 *Staphylococcus aureus*. *Mol Microbiol* 1995;16:895-907

1775 40. Ni Eidhin D, Perkins S, Francois P, Vaudaux P, Hook M and Foster TJ. Clumping
1776 factor B (ClfB), a new surface-located fibrinogen-binding adhesin of *Staphylococcus*
1777 *aureus*. *Mol Microbiol* 1998;30:245-57

1778 41. Mack D, Davies AP, Harris LG, Rohde H, Horstkotte MA and Knobloch JK.
1779 Microbial interactions in *Staphylococcus epidermidis* biofilms. *Anal Bioanal Chem*
1780 2007;387:399-408
1781 42. von Eiff C, Peters G and Heilmann C. Pathogenesis of infections due to coagulase-
1782 negative staphylococci. *Lancet Infect Dis* 2002;2:677-85
1783 43. Zhu Y, Weiss EC, Otto M, Fey PD, Smeltzer MS and Somerville GA.
1784 *Staphylococcus aureus* metabolism in a biofilm: the influence of arginine on
1785 polysaccharide intercellular adhesin synthesis, biofilm formation, and pathogenesis.
1786 *Infect Immun* 2007
1787 44. Murga R, Miller JM and Donlan RM. Biofilm formation by gram-negative bacteria
1788 on central venous catheter connectors: effect of conditioning films in a laboratory model.
1789 *J Clin Microbiol* 2001;39:2294-7
1790 45. Douglas LJ. *Candida* biofilms and their role in infection. *Trends Microbiol*
1791 2003;11:30-6
1792 46. Donlan RM. Biofilms: microbial life on surfaces. *Emerg Infect Dis* 2002;8:881-90
1793 47. Dunne WM, Jr., Burd EM. The effects of magnesium, calcium, EDTA, and pH on the
1794 in vitro adhesion of *Staphylococcus epidermidis* to plastic. *Microbiol Immunol*
1795 1992;36:1019-27
1796 48. Ozerdem Akpolat N, Elci S, Atmaca S, Akbayin H and Gul K. The effects of
1797 magnesium, calcium and EDTA on slime production by *Staphylococcus epidermidis*
1798 strains. *Folia Microbiol (Praha)* 2003;48:649-53
1799 49. Banin E, Brady KM and Greenberg EP. Chelator-induced dispersal and killing of
1800 *Pseudomonas aeruginosa* cells in a biofilm. *Appl Environ Microbiol* 2006;72:2064-9

Formatted: French (France)

1801 50. Donlan RM. Role of biofilms in antimicrobial resistance. *Asaio J* 2000;46:S47-52

1802 51. Farber BF, Kaplan MH and Clogston AG. *Staphylococcus epidermidis* extracted

1803 slime inhibits the antimicrobial action of glycopeptide antibiotics. *J Infect Dis*

1804 1990;161:37-40

1805 52. Branchini ML, Pfaller MA, Rhine-Chalberg J, Frempong T and Isenberg HD.

1806 Genotypic variation and slime production among blood and catheter isolates of *Candida*

1807 *parapsilosis*. *J Clin Microbiol* 1994;32:452-6

1808 53. Yoo S, Ha M, Choi D and Pai H. Effectiveness of surveillance of central catheter-

1809 related bloodstream infection in an ICU in Korea. *Infect Control Hosp Epidemiol*

1810 2001;22:433-6

1811 54. Warren DK, Zack JE, Cox MJ, Cohen MM and Fraser VJ. An educational

1812 intervention to prevent catheter-associated bloodstream infections in a non-teaching

1813 community medical center. *Crit Care Med* 2003;31

1814 55. Warren DK, Zack JE, Mayfield JL, et al. The effect of an education program on the

1815 incidence of central venous catheter-associated bloodstream infection in a medical ICU.

1816 *Chest* 2004;126:1612-8

1817 56. Warren DK, Cosgrove SE, Diekema DJ, et al. A multicenter intervention to prevent

1818 catheter-associated bloodstream infections. *Infect Control Hosp Epidemiol* 2006;27:662-

1819 9

1820 57. Higuera F, Rosenthal VD, Duarte P, Ruiz J, Franco G and Safdar N. The effect of

1821 process control on the incidence of central venous catheter-associated bloodstream

1822 infections and mortality in intensive care units in Mexico. *Crit Care Med* 2005;33:2022-7

Formatted: German (Germany)

Formatted: Italian (Italy)

1823 58. Coopersmith CM, Rebmann TL, Zack JE, et al. Effect of an education program on
 1824 decreasing catheter-related bloodstream infections in the surgical intensive care unit. Crit
 1825 Care Med 2002;30:59-64

1826 59. Coopersmith CM, Zack JE, Ward MR, et al. The impact of bedside behavior on
 1827 catheter-related bacteremia in the intensive care unit. Arch Surg 2004;139:131-6

1828 60. Sherertz RJ, Ely EW, Westbrook DM, et al. Education of physicians-in-training can
 1829 decrease the risk for vascular catheter infection. Ann Intern Med 2000;132:641-8

1830 61. Eggimann P, Harbarth S, Constantin MN, Touveneau S, Chevrolet JC and Pittet D.
 1831 Impact of a prevention strategy targeted at vascular-access care on incidence of infections
 1832 acquired in intensive care. Lancet 2000;355:1864-8

1833 62. Nehme AE. Nutritional support of the hospitalized patient. The team concept. JAMA
 1834 1980;243:1906-8

1835 63. Soifer NE, Borzak S, Edlin BR and Weinstein RA. Prevention of peripheral venous
 1836 catheter complications with an intravenous therapy team: a randomized controlled trial.
 1837 Arch Intern Med 1998;158:473-7

1838 64. Tomford JW, Hershey CO, McLaren CE, Porter DK and Cohen DI. Intravenous
 1839 therapy team and peripheral venous catheter-associated complications. A prospective
 1840 controlled study. Arch Intern Med 1984;144:1191-4

1841 65. Scalley RD, Van CS and Cochran RS. The impact of an i.v. team on the occurrence of
 1842 intravenous-related phlebitis. A 30-month study. J Intraven Nurs 1992;15:100-9

1843 66. Palefski SS, Stoddard GJ. The infusion nurse and patient complication rates of
 1844 peripheral-short catheters. A prospective evaluation. J Intraven Nurs 2001;24:113-23

Formatted: Swedish (Sweden)

1845 67. Miller JM, Goetz AM, Squier C and Muder RR. Reduction in nosocomial intravenous
1846 device-related bacteremias after institution of an intravenous therapy team. *J Intraven*
1847 *Nurs* 1996;19:103-6

1848 68. Hunter MR. Development of a Vascular Access Team in an acute care setting. *J Infus*
1849 *Nurs* 2003;26:86-91

1850 69. Hawes ML. A proactive approach to combating venous depletion in the hospital
1851 setting. *J Infus Nurs* 2007;30:33-44

1852 70. Brunelle D. Impact of a dedicated infusion therapy team on the reduction of catheter-
1853 related nosocomial infections. *J Infus Nurs* 2003;26:362-6

1854 71. Bosma TL, Jewesson PJ. An infusion program resource nurse consult service: our
1855 experience in a major Canadian teaching hospital. *J Infus Nurs* 2002;25:310-5

1856 72. Pierce CA, Baker JJ. A nursing process model: quantifying infusion therapy resource
1857 consumption. *J Infus Nurs* 2004;27:232-44

1858 73. Tomford JW, Hershey CO. The i.v. therapy team: impact on patient care and costs of
1859 hospitalization. *NITA* 1985;8:387-9

1860 74. Davis D, O'Brien MA, Freemantle N, Wolf FM, Mazmanian P and Taylor-Vaisey A.
1861 Impact of formal continuing medical education: do conferences, workshops, rounds, and
1862 other traditional continuing education activities change physician behavior or health care
1863 outcomes? *JAMA* 1999;282:867-74

1864 75. Alonso-Echanove J, Edwards JR, Richards MJ, et al. Effect of nurse staffing and
1865 antimicrobial-impregnated central venous catheters on the risk for bloodstream infections
1866 in intensive care units. *Infect Control Hosp Epidemiol* 2003;24:916-25

1867 76. Fridkin SK, Pear SM, Williamson TH, Galgiani JN and Jarvis WR. The role of
 1868 understaffing in central venous catheter-associated bloodstream infections. *Infect Control*
 1869 *Hosp Epidemiol* 1996;17:150-8

1870 77. Robert J, Fridkin SK, Blumberg HM, et al. The influence of the composition of the
 1871 nursing staff on primary bloodstream infection rates in a surgical intensive care unit.
 1872 *Infect Control Hosp Epidemiol* 2000;21:12-7.

1873 78. Sanders RA, Sheldon GF. Septic complications of total parenteral nutrition. A five
 1874 year experience. *Am J Surg* 1976;132:214-20

1875 79. Ryan JA, Jr., Abel RM, Abbott WM, et al. Catheter complications in total parenteral
 1876 nutrition. A prospective study of 200 consecutive patients. *N Engl J Med* 1974;290:757-
 1877 61

1878 80. Murphy LM, Lipman TO. Central venous catheter care in parenteral nutrition: a
 1879 review. *JPEN J Parenter Enteral Nutr* 1987;11:190-201

1880 81. Armstrong CW, Mayhall CG, Miller KB, et al. Prospective study of catheter
 1881 replacement and other risk factors for infection of hyperalimentation catheters. *J Infect*
 1882 *Dis* 1986;154:808-16

1883 82. Maki DG, Goldman DA and Rhame FS. Infection control in intravenous therapy. *Ann*
 1884 *Intern Med* 1973;79:867-87

1885 83. Band JD, Maki DG. Steel needles used for intravenous therapy. Morbidity in patients
 1886 with hematologic malignancy. *Arch Intern Med* 1980;140:31-4

1887 84. Tully JL, Friedland GH, Baldini LM and Goldmann DA. Complications of
 1888 intravenous therapy with steel needles and Teflon catheters. A comparative study. *Am J*
 1889 *Med* 1981;70:702-6

- 1890 85. Ryder MA. Peripheral access options. *Surg Oncol Clin N Am* 1995;4:395-427
- 1891 86. Parienti JJ, Thirion M, Megarbane B, et al. Femoral vs jugular venous catheterization
1892 and risk of nosocomial events in adults requiring acute renal replacement therapy: a
1893 randomized controlled trial. *JAMA* 2008;299:2413-22
- 1894 87. Moretti EW, Ofstead CL, Kristy RM and Wetzler HP. Impact of central venous
1895 catheter type and methods on catheter-related colonization and bacteraemia. *J Hosp Infect*
1896 2005;61:139-45
- 1897 88. Nagashima G, Kikuchi T, Tsuyuzaki H, et al. To reduce catheter-related bloodstream
1898 infections: is the subclavian route better than the jugular route for central venous
1899 catheterization? *J Infect Chemother* 2006;12:363-5
- 1900 89. Ruesch S, Walder B and Tramer MR. Complications of central venous catheters:
1901 internal jugular versus subclavian access--a systematic review. *Crit Care Med*
1902 2002;30:454-60
- 1903 90. Sadoyama G, Gontijo Filho PP. Comparison between the jugular and subclavian vein
1904 as insertion site for central venous catheters: microbiological aspects and risk factors for
1905 colonization and infection. *Braz J Infect Dis* 2003;7:142-8
- 1906 91. Heard SO, Wagle M, Vijayakumar E, et al. Influence of triple-lumen central venous
1907 catheters coated with chlorhexidine and silver sulfadiazine on the incidence of catheter-
1908 related bacteremia. *Arch Intern Med* 1998;158:81-7
- 1909 92. Richet H, Hubert B, Nitemberg G, et al. Prospective multicenter study of vascular-
1910 catheter-related complications and risk factors for positive central-catheter cultures in
1911 intensive care unit patients. *J Clin Microbiol* 1990;28:2520-5

1912 93. Safdar N, Kluger DM and Maki DG. A review of risk factors for catheter-related
1913 bloodstream infection caused by percutaneously inserted, noncuffed central venous
1914 catheters: implications for preventive strategies. *Medicine (Baltimore)* 2002;81:466-79
1915 94. Lorente L, Jimenez A, Iribarren JL, Jimenez JJ, Martin MM and Mora ML. The
1916 micro-organism responsible for central venous catheter-related bloodstream infection
1917 depends on catheter site. *Intensive Care Med* 2006;32:1449-50
1918 95. Traore O, Liotier J and Souweine B. Prospective study of arterial and central venous
1919 catheter colonization and of arterial- and central venous catheter-related bacteremia in
1920 intensive care units. *Crit Care Med* 2005;33:1276-80
1921 96. Joynt GM, Kew J, Gomersall CD, Leung VY and Liu EK. Deep venous thrombosis
1922 caused by femoral venous catheters in critically ill adult patients. *Chest* 2000;117:178-83.
1923 97. Mian NZ, Bayly R, Schreck DM, Besserman EB and Richmand D. Incidence of deep
1924 venous thrombosis associated with femoral venous catheterization. *Acad Emerg Med*
1925 1997;4:1118-21.
1926 98. Merrer J, De Jonghe B, Golliot F, et al. Complications of femoral and subclavian
1927 venous catheterization in critically ill patients: a randomized controlled trial. *JAMA*
1928 2001;286:700-7.
1929 99. Goetz AM, Wagener MM, Miller JM and Muder RR. Risk of infection due to central
1930 venous catheters: effect of site of placement and catheter type. *Infect Control Hosp*
1931 *Epidemiol* 1998;19:842-5
1932 100. Robinson JF, Robinson WA, Cohn A, Garg K and Armstrong JD, 2nd. Perforation
1933 of the great vessels during central venous line placement. *Arch Intern Med*
1934 1995;155:1225-8

Formatted: Italian (Italy)

Formatted: Swedish (Sweden)

1935 101. Trottier SJ, Veremakis C, O'Brien J and Auer AI. Femoral deep vein thrombosis
 1936 associated with central venous catheterization: results from a prospective, randomized
 1937 trial. Crit Care Med 1995;23:52-9

1938 102. Schillinger F, Schillinger D, Montagnac R and Milcent T. Post catheterisation vein
 1939 stenosis in haemodialysis: comparative angiographic study of 50 subclavian and 50
 1940 internal jugular accesses. Nephrol Dial Transplant 1991;6:722-4

1941 103. Cimochoowski GE, Worley E, Rutherford WE, Sartain J, Blondin J and Harter H.
 1942 Superiority of the internal jugular over the subclavian access for temporary dialysis.
 1943 Nephron 1990;54:154-61

1944 104. Barrett N, Spencer S, McIvor J and Brown EA. Subclavian stenosis: a major
 1945 complication of subclavian dialysis catheters. Nephrol Dial Transplant 1988;3:423-5

1946 105. Trerotola SO, Kuhn-Fulton J, Johnson MS, Shah H, Ambrosius WT and Kneebone
 1947 PH. Tunneled infusion catheters: increased incidence of symptomatic venous thrombosis
 1948 after subclavian versus internal jugular venous access. Radiology 2000;217:89-93

1949 106. Hind D, Calvert N, McWilliams R, et al. Ultrasonic locating devices for central
 1950 venous cannulation: meta-analysis. Bmj 2003;327:361

1951 107. Randolph AG, Cook DJ, Gonzales CA and Pribble CG. Ultrasound guidance for
 1952 placement of central venous catheters: a meta-analysis of the literature. Crit Care Med
 1953 1996;24:2053-8.

1954 108. Pronovost P, Needham D, Berenholtz S, et al. An intervention to decrease catheter-
 1955 related bloodstream infections in the ICU. N Engl J Med 2006;355:2725-32

1956 109. Berenholtz SM, Pronovost PJ, Lipsett PA, et al. Eliminating catheter-related
 1957 bloodstream infections in the intensive care unit. Crit Care Med 2004;32:2014-20

Formatted: Swedish (Sweden)

1958 110. Bansmer G, Keith D and Tesluk H. Complications following use of indwelling
 1959 catheters of inferior vena cava. JAMA 1958;167:1606-11

1960 111. Crane C. Venous interruption of septic thrombophlebitis. N Engl J Med
 1961 1960;262:947-51

1962 112. Indar R. The dangers of indwelling polyethelene cannulae in deep veins. Lancet
 1963 1959;1:284-6

1964 113. Maki DG, Mermel LA. Infections due to infusion therapy. In: JV B, PS B, eds.
 1965 Hospital Infections. 4 ed. Philadelphia: Lippencott-Raven, 1998:689-724

1966 114. Garland JS, Dunne WM, Jr., Havens P, et al. Peripheral intravenous catheter
 1967 complications in critically ill children: a prospective study. Pediatrics 1992;89:1145-50

1968 115. Garland JS, Nelson DB, Cheah TE, Hennes HH and Johnson TM. Infectious
 1969 complications during peripheral intravenous therapy with Teflon catheters: a prospective
 1970 study. Pediatr Infect Dis J 1987;6:918-21

1971 116. Breschan C, Platzer M, Jost R, Schaumberger F, Stettner H and Likar R.
 1972 Comparison of catheter-related infection and tip colonization between internal jugular
 1973 and subclavian central venous catheters in surgical neonates. Anesthesiology
 1974 2007;107:946-53

1975 117. Deshpande KS, Hatem C, Ulrich HL, et al. The incidence of infectious
 1976 complications of central venous catheters at the subclavian, internal jugular, and femoral
 1977 sites in an intensive care unit population. Crit Care Med 2005;33:13-20; discussion 234-5

1978 118. Durbec O, Viviand X, Potie F, Vialet R, Albanese J and Martin C. A prospective
 1979 evaluation of the use of femoral venous catheters in critically ill adults. Crit Care Med
 1980 1997;25:1986-9.

Formatted: Swedish (Sweden)

Formatted: German (Germany)

1981 119. Venkataraman ST, Thompson AE and Orr RA. Femoral vascular catheterization in
 1982 critically ill infants and children. Clin Pediatr (Phila) 1997;36:311-9.

1983 120. Sheridan RL, Weber JM. Mechanical and infectious complications of central venous
 1984 cannulation in children: lessons learned from a 10-year experience placing more than
 1985 1000 catheters. J Burn Care Res 2006;27:713-8

1986 121. Stenzel JP, Green TP, Fuhrman BP, Carlson PE and Marchessault RP. Percutaneous
 1987 central venous catheterization in a pediatric intensive care unit: a survival analysis of
 1988 complications. Crit Care Med 1989;17:984-8

1989 122. Goldstein AM, Weber JM and Sheridan RL. Femoral venous access is safe in
 1990 burned children: an analysis of 224 catheters. J Pediatr 1997;130:442-6.

1991 123. Ramos GE, Bolgiani AN, Patino O, et al. Catheter infection risk related to the
 1992 distance between insertion site and burned area. J Burn Care Rehabil 2002;23:266-71

1993 124. Sheth NK, Franson TR, Rose HD, Buckmire FL, Cooper JA and Sohnle PG.
 1994 Colonization of bacteria on polyvinyl chloride and Teflon intravascular catheters in
 1995 hospitalized patients. J Clin Microbiol 1983;18:1061-3

1996 125. Maki DG, Ringer M. Risk factors for infusion-related phlebitis with small peripheral
 1997 venous catheters. A randomized controlled trial. Ann Intern Med 1991;114:845-54

1998 126. Maki DG, Ringer M. Evaluation of dressing regimens for prevention of infection
 1999 with peripheral intravenous catheters. Gauze, a transparent polyurethane dressing, and an
 2000 iodophor-transparent dressing. JAMA 1987;258:2396-403

2001 127. Boyce JM, Pittet D. Guideline for Hand Hygiene in Health-Care Settings:
 2002 recommendations of the Healthcare Infection Control Practices Advisory Committee and

Formatted: Portuguese (Brazil)

2003 the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Infect Control Hosp

2004 Epidemiol 2002;23:S3-40

2005 128. Bischoff WE, Reynolds TM, Sessler CN, Edmond MB and Wenzel RP.

2006 Handwashing compliance by healthcare workers: The impact of introducing an

2007 accessible, alcohol-based hand antiseptic. Arch Intern Med 2000;160:1017-21

2008 129. Boyce JM, Kelliher S and Vallande N. Skin irritation and dryness associated with

2009 two hand-hygiene regimens: soap-and-water hand washing versus hand antisepsis with an

2010 alcoholic hand gel. Infect Control Hosp Epidemiol 2000;21:442-8.

2011 130. Pittet D, Dharan S, Touveneau S, Sauvan V and Perneger TV. Bacterial

2012 contamination of the hands of hospital staff during routine patient care. Arch Intern Med

2013 1999;159:821-6

2014 131. Simmons B, Bryant J, Neiman K, Spencer L and Arheart K. The role of

2015 handwashing in prevention of endemic intensive care unit infections. Infect Control Hosp

2016 Epidemiol 1990;11:589-94

2017 132. Raad, II, Hohn DC, Gilbreath BJ, et al. Prevention of central venous catheter-related

2018 infections by using maximal sterile barrier precautions during insertion. Infect Control

2019 Hosp Epidemiol 1994;15:231-8

2020 133. Capdevila JA, Segarra A and Pahissa A. Catheter-related bacteremia in patients

2021 undergoing hemodialysis. Ann Intern Med 1998;128:600

Formatted: Swedish (Sweden)

2022 134. Abi-Said D, Raad I, Umphrey J, et al. Infusion therapy team and dressing changes of

2023 central venous catheters. Infect Control Hosp Epidemiol 1999;20:101-5

2024 135. Pittet D, Hugonnet S, Harbath S, et al. Effectiveness of a hospital-wide programme to

2025 improve compliance with hand hygiene. Lancet 2000;356:1307-9

Formatted: French (France)

2026 136. Carrer S, Bocchi A, Bortolotti M, et al. Effect of different sterile barrier precautions
2027 and central venous catheter dressing on the skin colonization around the insertion site.

2028 Minerva Anestesiol 2005;71:197-206

Formatted: Italian (Italy)

2029 137. Mermel LA, Maki DG. Infectious complications of Swan-Ganz pulmonary artery
2030 catheters. Pathogenesis, epidemiology, prevention, and management. Am J Respir Crit

2031 Care Med 1994;149:1020-36

2032 138. Cohen Y, Fosse JP, Karoubi P, et al. The "hands-off" catheter and the prevention of
2033 systemic infections associated with pulmonary artery catheter: a prospective study. Am J
2034 Respir Crit Care Med 1998;157:284-7

Formatted: French (France)

2035 139. Maki DG, Band JD. A comparative study of polyantibiotic and iodophor ointments
2036 in prevention of vascular catheter-related infection. Am J Med 1981;70:739-44

Formatted: Swedish (Sweden)

2037 140. Maki DG, Ringer M and Alvarado CJ. Prospective randomised trial of povidone-
2038 iodine, alcohol, and chlorhexidine for prevention of infection associated with central
2039 venous and arterial catheters. Lancet 1991;338:339-43

2040 141. Mimos O, Pieroni L, Lawrence C, et al. Prospective, randomized trial of two

Formatted: French (France)

2041 antiseptic solutions for prevention of central venous or arterial catheter colonization and
2042 infection in intensive care unit patients. Crit Care Med 1996;24:1818-23

2043 142. Humar A, Ostromecki A, Drenfeld J, et al. Prospective randomized trial of 10%
2044 povidone-iodine versus 0.5% tincture of chlorhexidine as cutaneous antisepsis for
2045 prevention of central venous catheter infection. Clin Infect Dis 2000;31:1001-7

2046 143. Chaiyakunapruk N, Veenstra DL, Lipsky BA and Saint S. Chlorhexidine compared
2047 with povidone-iodine solution for vascular catheter-site care: a meta-analysis. Ann Intern
2048 Med 2002;136:792-801

2049 144. Chaiyakunapruk N, Veenstra DL, Lipsky BA, Sullivan SD and Saint S. Vascular
 2050 catheter site care: the clinical and economic benefits of chlorhexidine gluconate
 2051 compared with povidone iodine. Clin Infect Dis 2003;37:764-71

2052 145. Parienti JJ, du Cheyron D, Ramakers M, et al. Alcoholic povidone-iodine to prevent
 2053 central venous catheter colonization: A randomized unit-crossover study. Crit Care Med
 2054 2004;32:708-13

2055 146. Maki DG, Stolz SS, Wheeler S and Mermel LA. A prospective, randomized trial of
 2056 gauze and two polyurethane dressings for site care of pulmonary artery catheters:
 2057 implications for catheter management. Crit Care Med 1994;22:1729-37

2058 147. Bijma R, Girbes AR, Kleijer DJ and Zwaveling JH. Preventing central venous
 2059 catheter-related infection in a surgical intensive-care unit. Infect Control Hosp Epidemiol
 2060 1999;20:618-20

2061 148. Madeo M, Martin CR, Turner C, Kirkby V and Thompson DR. A randomized trial
 2062 comparing Arglaes (a transparent dressing containing silver ions) to Tegaderm (a
 2063 transparent polyurethane dressing) for dressing peripheral arterial catheters and central
 2064 vascular catheters. Intensive Crit Care Nurs 1998;14:187-91

2065 149. Laura R, Degl'Innocenti M, Mocali M, et al. Comparison of two different time
 2066 interval protocols for central venous catheter dressing in bone marrow transplant patients:
 2067 results of a randomized, multicenter study. The Italian Nurse Bone Marrow Transplant
 2068 Group (GITMO). Haematologica 2000;85:275-9

2069 150. Zakrzewska-Bode A, Muyltjens HL, Liem KD and Hoogkamp-Korstanje JA.
 2070 Mupirocin resistance in coagulase-negative staphylococci, after topical prophylaxis for
 2071 the reduction of colonization of central venous catheters. J Hosp Infect 1995;31:189-93

Formatted: French (France)

Formatted: French (France)

2072 151. Flowers RH, Schwenzer KJ, Kopel RF, Fisch MJ, Tucker SI and Farr BM. Efficacy
 2073 of an attachable subcutaneous cuff for the prevention of intravascular catheter-related
 2074 infection. A randomized, controlled trial. JAMA 1989;261:878-83

2075 152. Robbins J, Cromwell P and Korones DN. Swimming and central venous catheter-
 2076 related infections in the child with cancer. J Pediatr Oncol Nurs 1999;16:51-6

2077 153. Howell PB, Walters PE, Donowitz GR and Farr BM. Risk factors for infection of
 2078 adult patients with cancer who have tunnelled central venous catheters. Cancer
 2079 1995;75:1367-75

2080 154. Rao SP, Oreopoulos DG. Unusual complications of a polyurethane PD catheter.
 2081 Perit Dial Int 1997;17:410-2

2082 155. Riu S, Ruiz CG, Martinez-Vea A, Peralta C and Oliver JA. Spontaneous rupture of
 2083 polyurethane peritoneal catheter. A possible deleterious effect of mupirocin ointment.
 2084 Nephrol Dial Transplant 1998;13:1870-1

2085 156. Timsit JF, Schwebel C, Bouadma L, et al. Chlorhexidine-impregnated sponges and
 2086 less frequent dressing changes for prevention of catheter-related infections in critically ill
 2087 adults: a randomized controlled trial. JAMA 2009;301:1231-41

2088 157. Ho KM, Litton E. Use of chlorhexidine-impregnated dressing to prevent vascular
 2089 and epidural catheter colonization and infection: a meta-analysis. J Antimicrob
 2090 Chemother 2006;58:281-7

2091 158. Levy I, Katz J, Solter E, et al. Chlorhexidine-impregnated dressing for prevention of
 2092 colonization of central venous catheters in infants and children: a randomized controlled
 2093 study. Pediatr Infect Dis J 2005;24:676-9

Formatted: Italian (Italy)

2094 159. Hoffmann KK, Weber DJ, Samsa GP and Rutala WA. Transparent polyurethane
 2095 film as an intravenous catheter dressing. A meta-analysis of the infection risks. JAMA
 2096 1992;267:2072-6

2097 160. Gillies D, O'Riordan E, Carr D, O'Brien I, Frost J and Gunning R. Central venous
 2098 catheter dressings: a systematic review. J Adv Nurs 2003;44:623-32

2099 161. Ruschulte H, Franke M, Gastmeier P, et al. Prevention of central venous catheter-
 2100 related infections with chlorhexidine gluconate impregnated wound dressings: a
 2101 randomized controlled trial. Ann Hematol 2009;88:267-72

2102 162. Bleasdale SC, Trick WE, Gonzalez IM, Lyles RD, Hayden MK and Weinstein RA.
 2103 Effectiveness of chlorhexidine bathing to reduce catheter-associated bloodstream
 2104 infections in medical intensive care unit patients. Arch Intern Med 2007;167:2073-9

2105 163. Yamamoto AJ, Solomon JA, Soulen MC, et al. Sutureless securement device
 2106 reduces complications of peripherally inserted central venous catheters. J Vasc Interv
 2107 Radiol 2002;13:77-81

2108 164. Raad I, Darouiche R, Dupuis J, et al. Central venous catheters coated with
 2109 minocycline and rifampin for the prevention of catheter-related colonization and
 2110 bloodstream infections. A randomized, double-blind trial. The Texas Medical Center
 2111 Catheter Study Group. Ann Intern Med 1997;127:267-74

2112 165. Bhutta A, Gilliam C, Honeycutt M, et al. Reduction of bloodstream infections
 2113 associated with catheters in paediatric intensive care unit: stepwise approach. Bmj
 2114 2007;334:362-5

Formatted: German (Germany)

Formatted: Swedish (Sweden)

Formatted: Italian (Italy)

Formatted: Swedish (Sweden)

2115 166. Chelliah A, Heydon KH, Zaoutis TE, et al. Observational trial of antibiotic-coated
2116 central venous catheters in critically ill pediatric patients. *Pediatr Infect Dis J*
2117 2007;26:816-20

2118 167. Veenstra DL, Saint S, Saha S, Lumley T and Sullivan SD. Efficacy of antiseptic-
2119 impregnated central venous catheters in preventing catheter-related bloodstream
2120 infection: a meta-analysis. *JAMA* 1999;281:261-7

2121 168. Maki DG, Stolz SM, Wheeler S and Mermel LA. Prevention of central venous
2122 catheter-related bloodstream infection by use of an antiseptic-impregnated catheter. A
2123 randomized, controlled trial. *Ann Intern Med* 1997;127:257-66

2124 169. Brun-Buisson C, Doyon F, Sollet JP, Cochard JF, Cohen Y and Nitenberg G.
2125 Prevention of intravascular catheter-related infection with newer chlorhexidine-silver
2126 sulfadiazine-coated catheters: a randomized controlled trial. *Intensive Care Med*
2127 2004;30:837-43

2128 170. Ostendorf T, Meinhold A, Harter C, et al. Chlorhexidine and silver-sulfadiazine
2129 coated central venous catheters in haematological patients--a double-blind, randomised,
2130 prospective, controlled trial. *Support Care Cancer* 2005;13:993-1000

2131 171. Rupp ME, Lisco SJ, Lipsett PA, et al. Effect of a second-generation venous catheter
2132 impregnated with chlorhexidine and silver sulfadiazine on central catheter-related
2133 infections: a randomized, controlled trial. *Ann Intern Med* 2005;143:570-80

2134 172. Bassetti S, Hu J, D'Agostino RB, Jr. and Sherertz RJ. Prolonged antimicrobial
2135 activity of a catheter containing chlorhexidine-silver sulfadiazine extends protection
2136 against catheter infections in vivo. *Antimicrob Agents Chemother* 2001;45:1535-8.

2137 173. Oda T, Hamasaki J, Kanda N and Mikami K. Anaphylactic shock induced by an
 2138 antiseptic-coated central venous [correction of nervous] catheter. Anesthesiology
 2139 1997;87:1242-4.

2140 174. Pittaway A, Ford S. Allergy to chlorhexidine-coated central venous catheters
 2141 revisited. Br J Anaesth 2002;88:304-5; author reply 305

2142 175. Stephens R, Mythen M, Kallis P, Davies DW, Egner W and Rickards A. Two
 2143 episodes of life-threatening anaphylaxis in the same patient to a chlorhexidine-
 2144 sulphadiazine-coated central venous catheter. Br J Anaesth 2001;87:306-8

2145 176. Terazawa E, Shimonaka H, Nagase K, Masue T and Dohi S. Severe anaphylactic
 2146 reaction due to a chlorhexidine-impregnated central venous catheter. Anesthesiology
 2147 1998;89:1296-8

2148 177. Veenstra DL, Saint S and Sullivan SD. Cost-effectiveness of antiseptic-impregnated
 2149 central venous catheters for the prevention of catheter-related bloodstream infection.
 2150 JAMA 1999;282:554-60

2151 178. Darouiche RO, Raad, II, Heard SO, et al. A comparison of two antimicrobial-
 2152 impregnated central venous catheters. Catheter Study Group. N Engl J Med 1999;340:1-8

2153 179. Hanna H, Benjamin R, Chatzinikolaou I, et al. Long-term silicone central venous
 2154 catheters impregnated with minocycline and rifampin decrease rates of catheter-related
 2155 bloodstream infection in cancer patients: a prospective randomized clinical trial. J Clin
 2156 Oncol 2004;22:3163-71

2157 180. Tambe SM, Sampath L and Modak SM. In vitro evaluation of the risk of developing
 2158 bacterial resistance to antiseptics and antibiotics used in medical devices. J Antimicrob
 2159 Chemother 2001;47:589-98

Formatted: Swedish (Sweden)

2160 181. Sampath LA, Tambe SM and Modak SM. In vitro and in vivo efficacy of catheters
 2161 impregnated with antiseptics or antibiotics: evaluation of the risk of bacterial resistance to
 2162 the antimicrobials in the catheters. Infect Control Hosp Epidemiol 2001;22:640-6

2163 182. Marciante KD, Veenstra DL, Lipsky BA and Saint S. Which antimicrobial
 2164 impregnated central venous catheter should we use? Modeling the costs and outcomes of
 2165 antimicrobial catheter use. Am J Infect Control 2003;31:1-8

2166 183. Shorr AF, Humphreys CW and Helman DL. New choices for central venous
 2167 catheters: potential financial implications. Chest 2003;124:275-84

2168 184. Hagau N, Studnicska D, Gavrus RL, Csipak G, Hagau R and Slavcovici AV. Central
 2169 venous catheter colonization and catheter-related bloodstream infections in critically ill
 2170 patients: a comparison between standard and silver-integrated catheters. Eur J
 2171 Anaesthesiol 2009;26:752-8

2172 185. Bong JJ, Kite P, Wilco MH and McMahon MJ. Prevention of catheter-related
 2173 bloodstream infection by silver iontophoretic central venous catheters: a randomised
 2174 controlled trial. J Clin Pathol 2003;56:731-5

2175 186. Corral L, Nolla-Salas M, Ibanez-Nolla J, et al. A prospective, randomized study in
 2176 critically ill patients using the Oligon Vantex catheter. J Hosp Infect 2003;55:212-9

2177 187. Ranucci M, Isgro G, Giomarelli PP, et al. Impact of oligon central venous catheters
 2178 on catheter colonization and catheter-related bloodstream infection. Crit Care Med
 2179 2003;31:52-9

2180 188. van de Wetering MD, van Woensel JB. Prophylactic antibiotics for preventing early
 2181 central venous catheter Gram positive infections in oncology patients. Cochrane Database
 2182 Syst Rev 2007:CD003295

Formatted: French (France)

Formatted: Swedish (Sweden)

2183 189. Raad, II, Hachem RY, Abi-Said D, et al. A prospective crossover randomized trial
 2184 of novobiocin and rifampin prophylaxis for the prevention of intravascular catheter
 2185 infections in cancer patients treated with interleukin-2. *Cancer* 1998;82:403-11

2186 190. McKee R, Dunsmuir R, Whitby M and Garden OJ. Does antibiotic prophylaxis at
 2187 the time of catheter insertion reduce the incidence of catheter-related sepsis in
 2188 intravenous nutrition? *J Hosp Infect* 1985;6:419-25

2189 191. Sandoe JA, Kumar B, Stoddart B, et al. Effect of extended perioperative antibiotic
 2190 prophylaxis on intravascular catheter colonization and infection in cardiothoracic surgery
 2191 patients. *J Antimicrob Chemother* 2003;52:877-9

2192 192. Inglis GD, Jardine LA and Davies MW. Prophylactic antibiotics to reduce morbidity
 2193 and mortality in neonates with umbilical artery catheters. *Cochrane Database Syst Rev*
 2194 2007:CD004697

2195 193. Craft AP, Finan NN and Barrington KJ. Vancomycin for prophylaxis against sepsis
 2196 in preterm neonates. *Cochrane Database Syst Rev* 2000:CD001971

2197 194. Fukunaga A, Naritaka H, Fukaya R, Tabuse M and Nakamura T. Povidone-iodine
 2198 ointment and gauze dressings associated with reduced catheter-related infection in
 2199 seriously ill neurosurgical patients. *Infect Control Hosp Epidemiol* 2004;25:696-8

2200 195. Johnson DW, MacGinley R, Kay TD, et al. A randomized controlled trial of topical
 2201 exit site mupirocin application in patients with tunnelled, cuffed haemodialysis catheters.
 2202 *Nephrol Dial Transplant* 2002;17:1802-7

2203 196. Fong IW. Prevention of haemodialysis and peritoneal dialysis catheter-related
 2204 infection by topical povidone-iodine. *Postgrad Med J* 1993;69 Suppl 3:S15-7

Formatted: French (France)

Formatted: Swedish (Sweden)

Formatted: Portuguese (Brazil)

Formatted: Swedish (Sweden)

2205 197. Levin A, Mason AJ, Jindal KK, Fong IW and Goldstein MB. Prevention of
 2206 hemodialysis subclavian vein catheter infections by topical povidone-iodine. *Kidney Int*
 2207 1991;40:934-8

2208 198. National Kidney Foundation. III. NKF-K/DOQI Clinical Practice Guidelines for
 2209 Vascular Access: update 2000. *Am J Kidney Dis* 2001;37:S137-81.

2210 199. Norden CW. Application of antibiotic ointment to the site of venous catheterization-
 2211 -a controlled trial. *J Infect Dis* 1969;120:611-5.

2212 200. Zinner SH, Denny-Brown BC, Braun P, Burke JP, Toala P and Kass EH. Risk of
 2213 infection with intravenous indwelling catheters: effect of application of antibiotic
 2214 ointment. *J Infect Dis* 1969;120:616-9.

2215 201. von Eiff C, Becker K, Machka K, Stammer H and Peters G. Nasal Carriage as a
 2216 Source of *Staphylococcus aureus* Bacteremia. *N Engl J Med* 2001;344:11-16.

2217 202. Chow JW, Yu VL. *Staphylococcus aureus* nasal carriage in hemodialysis patients.
 2218 Its role in infection and approaches to prophylaxis. *Arch Intern Med* 1989;149:1258-62

2219 203. Yu VL, Goetz A, Wagener M, et al. *Staphylococcus aureus* nasal carriage and
 2220 infection in patients on hemodialysis. Efficacy of antibiotic prophylaxis. *N Engl J Med*
 2221 1986;315:91-6

2222 204. Casewell MW. The nose: an underestimated source of *Staphylococcus aureus*
 2223 causing wound infection. *J Hosp Infect* 1998;40:S3-11

2224 205. Hill RL, Fisher AP, Ware RJ, Wilson S and Casewell MW. Mupirocin for the
 2225 reduction of colonization of internal jugular cannulae--a randomized controlled trial. *J*
 2226 *Hosp Infect* 1990;15:311-21

Formatted: French (France)

2227 206. Sesso R, Barbosa D, Leme IL, et al. Staphylococcus aureus prophylaxis in
 2228 hemodialysis patients using central venous catheter: effect of mupirocin ointment. J Am
 2229 Soc Nephrol 1998;9:1085-92

2230 207. Boelaert JR, Van Landuyt HW, Godard CA, et al. Nasal mupirocin ointment
 2231 decreases the incidence of Staphylococcus aureus bacteraemias in haemodialysis patients.
 2232 Nephrol Dial Transplant 1993;8:235-9

2233 208. Netto dos Santos KR, de Souza Fonseca L and Gontijo Filho PP. Emergence of
 2234 high-level mupirocin resistance in methicillin-resistant Staphylococcus aureus isolated
 2235 from Brazilian university hospitals. Infect Control Hosp Epidemiol 1996;17:813-6

2236 209. Miller MA, Dascal A, Portnoy J and Mendelson J. Development of mupirocin
 2237 resistance among methicillin-resistant Staphylococcus aureus after widespread use of
 2238 nasal mupirocin ointment. Infect Control Hosp Epidemiol 1996;17:811-3

2239 210. Lok CE, Stanley KE, Hux JE, Richardson R, Tobe SW and Conly J. Hemodialysis
 2240 infection prevention with polysporin ointment. J Am Soc Nephrol 2003;14:169-79

2241 211. Schwartz C, Henrickson KJ, Roghmann K and Powell K. Prevention of bacteremia
 2242 attributed to luminal colonization of tunneled central venous catheters with vancomycin-
 2243 susceptible organisms. J Clin Oncol 1990;8:1591-7

2244 212. Rackoff WR, Weiman M, Jakobowski D, et al. A randomized, controlled trial of the
 2245 efficacy of a heparin and vancomycin solution in preventing central venous catheter
 2246 infections in children. J Pediatr 1995;127:147-51

2247 213. Carratala J, Niubo J, Fernandez-Sevilla A, et al. Randomized, double-blind trial of
 2248 an antibiotic-lock technique for prevention of gram-positive central venous catheter-

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

2249 related infection in neutropenic patients with cancer. Antimicrob Agents Chemother
 2250 1999;43:2200-4

2251 214. Jurewitsch B, Lee T, Park J and Jeejeebhoy K. Taurolidine 2% as an antimicrobial
 2252 lock solution for prevention of recurrent catheter-related bloodstream infections. J
 2253 Parenter Enteral Nutr 1998;22:242-4

2254 215. Henrickson KJ, Axtell RA, Hoover SM, et al. Prevention of central venous catheter-
 2255 related infections and thrombotic events in immunocompromised children by the use of
 2256 vancomycin/ciprofloxacin/heparin flush solution: A randomized, multicenter, double-
 2257 blind trial. J Clin Oncol 2000;18:1269-78

2258 216. Daghistani D, Horn M, Rodriguez Z, Schoenike S and Toledano S. Prevention of
 2259 indwelling central venous catheter sepsis. Med Pediatr Oncol 1996;26:405-8

2260 217. Barriga FJ, Varas M, Potin M, et al. Efficacy of a vancomycin solution to prevent
 2261 bacteremia associated with an indwelling central venous catheter in neutropenic and non-
 2262 neutropenic cancer patients. Med Pediatr Oncol 1997;28:196-200

2263 218. Dogra GK, Herson H, Hutchison B, et al. Prevention of tunneled hemodialysis
 2264 catheter-related infections using catheter-restricted filling with gentamicin and citrate: a
 2265 randomized controlled study. J Am Soc Nephrol 2002;13:2133-9

2266 219. Allon M. Prophylaxis against dialysis catheter-related bacteremia with a novel
 2267 antimicrobial lock solution. Clin Infect Dis 2003;36:1539-44

2268 220. Elhassan NO, Stevens TP, Gigliotti F, Hardy DJ, Cole CA and Sinkin RA.
 2269 Vancomycin usage in central venous catheters in a neonatal intensive care unit. Pediatr
 2270 Infect Dis J 2004;23:201-6

Formatted: Swedish (Sweden)

Formatted: Swedish (Sweden)

2271 221. McIntyre CW, Hulme LJ, Taal M and Fluck RJ. Locking of tunneled hemodialysis
2272 catheters with gentamicin and heparin. Kidney Int 2004;66:801-5

2273 222. Betjes MG, van Agteren M. Prevention of dialysis catheter-related sepsis with a
2274 citrate-taurolidine-containing lock solution. Nephrol Dial Transplant 2004;19:1546-51

Formatted: Swedish (Sweden)

2275 223. Weijmer MC, van den Dorpel MA, Van de Ven PJ, et al. Randomized, clinical trial
2276 comparison of trisodium citrate 30% and heparin as catheter-locking solution in

2277 hemodialysis patients. J Am Soc Nephrol 2005;16:2769-77

2278 224. Bleyer AJ, Mason L, Russell G, Raad, II and Sherertz RJ. A randomized, controlled
2279 trial of a new vascular catheter flush solution (minocycline-EDTA) in temporary

2280 hemodialysis access. Infect Control Hosp Epidemiol 2005;26:520-4

2281 225. Kim SH, Song KI, Chang JW, et al. Prevention of uncuffed hemodialysis catheter-
2282 related bacteremia using an antibiotic lock technique: a prospective, randomized clinical
2283 trial. Kidney Int 2006;69:161-4

Formatted: French (France)

2284 226. Al-Hwiesh AK, Abdul-Rahman IS. Successful prevention of tunneled, central
2285 catheter infection by antibiotic lock therapy using vancomycin and gentamycin. Saudi J
2286 Kidney Dis Transpl 2007;18:239-47

2287 227. Nori US, Manoharan A, Yee J and Besarab A. Comparison of low-dose gentamicin
2288 with minocycline as catheter lock solutions in the prevention of catheter-related

2289 bacteremia. Am J Kidney Dis 2006;48:596-605

2290 228. Saxena AK, Panhotra BR, Sundaram DS, et al. Tunneled catheters' outcome

Formatted: Swedish (Sweden)

2291 optimization among diabetics on dialysis through antibiotic-lock placement. Kidney Int
2292 2006;70:1629-35

2293 229. Yahav D, Rozen-Zvi B, Gaftor-Gvili A, Leibovici L, Gaftor U and Paul M.
 2294 Antimicrobial lock solutions for the prevention of infections associated with intravascular
 2295 catheters in patients undergoing hemodialysis: systematic review and meta-analysis of
 2296 randomized, controlled trials. Clin Infect Dis 2008;47:83-93

2297 230. Labriola L, Crott R and Jadoul M. Preventing haemodialysis catheter-related
 2298 bacteraemia with an antimicrobial lock solution: a meta-analysis of prospective
 2299 randomized trials. Nephrol Dial Transplant 2008;23:1666-72

2300 231. Jaffer Y, Selby NM, Taal MW, Fluck RJ and McIntyre CW. A meta-analysis of
 2301 hemodialysis catheter locking solutions in the prevention of catheter-related infection.
 2302 Am J Kidney Dis 2008;51:233-41

2303 232. Safdar N, Maki DG. Use of vancomycin-containing lock or flush solutions for
 2304 prevention of bloodstream infection associated with central venous access devices: a
 2305 meta-analysis of prospective, randomized trials. Clin Infect Dis 2006;43:474-84

2306 233. Sanders J, Pithie A, Ganly P, et al. A prospective double-blind randomized trial
 2307 comparing intraluminal ethanol with heparinized saline for the prevention of catheter-
 2308 associated bloodstream infection in immunosuppressed haematology patients. J
 2309 Antimicrob Chemother 2008;62:809-15

2310 234. Randolph AG, Cook DJ, Gonzales CA and Andrew M. Benefit of heparin in central
 2311 venous and pulmonary artery catheters: a meta-analysis of randomized controlled trials.
 2312 Chest 1998;113:165-71

2313 235. Schinabeck MK gM. Biofilm-Related Indwelling Medical Device Infections. In:
 2314 Pace JL RM, Finch RG, ed. Biofilms, Infection, and Antimicrobial Therapy. Boca Raton:
 2315 Taylor and Francis, 2006:39-50

2316 236. Gristina AG. Biomaterial-centered infection: microbial adhesion versus tissue
 2317 integration. *Science* 1987;237:1588-95

2318 237. Timsit JF, Farkas JC, Boyer JM, et al. Central vein catheter-related thrombosis in
 2319 intensive care patients: incidence, risks factors, and relationship with catheter-related
 2320 sepsis. *Chest* 1998;114:207-13

2321 238. Eastman ME, Khorsand M, Maki DG, et al. Central venous device-related infection
 2322 and thrombosis in patients treated with moderate dose continuous-infusion interleukin-2.
 2323 *Cancer* 2001;91:806-14

2324 239. Abdelkefi A, Torjman L, Ladeb S, et al. Randomized trial of prevention of catheter-
 2325 related bloodstream infection by continuous infusion of low-dose unfractionated heparin
 2326 in patients with hematologic and oncologic disease. *J Clin Oncol* 2005;23:7864-70

2327 240. Mermel LA, Stolz SM and Maki DG. Surface antimicrobial activity of heparin-
 2328 bonded and antiseptic-impregnated vascular catheters. *J Infect Dis* 1993;167:920-4.

2329 241. Pierce CM, Wade A and Mok Q. Heparin-bonded central venous lines reduce
 2330 thrombotic and infective complications in critically ill children. *Intensive Care Med*
 2331 2000;26:967-72.

2332 242. Appelgren P, Ransjo U, Bindslev L, Espersen F and Larm O. Surface heparinization
 2333 of central venous catheters reduces microbial colonization in vitro and in vivo: results
 2334 from a prospective, randomized trial. *Crit Care Med* 1996;24:1482-9

2335 243. Abdelkefi A, Achour W, Ben Othman T, et al. Use of heparin-coated central venous
 2336 lines to prevent catheter-related bloodstream infection. *J Support Oncol* 2007;5:273-8

Formatted: Italian (Italy)

2337 244. Carrasco MN, Bueno A, de las Cuevas C, et al. Evaluation of a triple-lumen central
2338 venous heparin-coated catheter versus a catheter coated with chlorhexidine and silver
2339 sulfadiazine in critically ill patients. Intensive Care Med 2004;30:633-8

2340 245. Levy JH, Hursting MJ. Heparin-induced thrombocytopenia, a prothrombotic disease.
2341 Hematol Oncol Clin North Am 2007;21:65-88

2342 246. Weijmer MC, Debets-Ossenkopp YJ, Van De Vondervoort FJ and ter Wee PM.
2343 Superior antimicrobial activity of trisodium citrate over heparin for catheter locking.
2344 Nephrol Dial Transplant 2002;17:2189-95

2345 247. Boraks P, Seale J, Price J, et al. Prevention of central venous catheter associated
2346 thrombosis using minidose warfarin in patients with haematological malignancies. Br J
2347 Haematol 1998;101:483-6

2348 248. Bern MM, Lokich JJ, Wallach SR, et al. Very low doses of warfarin can prevent
2349 thrombosis in central venous catheters. A randomized prospective trial. Ann Intern Med
2350 1990;112:423-8

2351 249. Akl EA, Karmath G, Yosucio V, et al. Anticoagulation for thrombosis prophylaxis
2352 in cancer patients with central venous catheters. Cochrane Database Syst Rev
2353 2007:CD006468

2354 250. Akl EA, Muti P and Schunemann HJ. Anticoagulation in patients with cancer: an
2355 overview of reviews. Pol Arch Med Wewn 2008;118:183-93

2356 251. Klerk CP, Smorenburg SM and Buller HR. Thrombosis prophylaxis in patient
2357 populations with a central venous catheter: a systematic review. Arch Intern Med
2358 2003;163:1913-21

Formatted: German (Germany)

2359 252. Heaton DC, Han DY and Inder A. Minidose (1 mg) warfarin as prophylaxis for
 2360 central vein catheter thrombosis. Intern Med J 2002;32:84-8

2361 253. Masci G, Magagnoli M, Zucali PA, et al. Minidose warfarin prophylaxis for
 2362 catheter-associated thrombosis in cancer patients: can it be safely associated with
 2363 fluorouracil-based chemotherapy? J Clin Oncol 2003;21:736-9

2364 254. Kuter DJ. Thrombotic complications of central venous catheters in cancer patients.
 2365 Oncologist 2004;9:207-16

2366 255. Maki DG, Botticelli JT, LeRoy ML and Thielke TS. Prospective study of replacing
 2367 administration sets for intravenous therapy at 48- vs 72-hour intervals. 72 hours is safe
 2368 and cost-effective. JAMA 1987;258:1777-81

2369 256. Collin J, Collin C. Infusion thrombophlebitis. Lancet 1975;2:458

2370 257. Lai KK. Safety of prolonging peripheral cannula and i.v. tubing use from 72 hours
 2371 to 96 hours. Am J Infect Control 1998;26:66-70

2372 258. Fontaine PJ. Performance of a new softening expanding midline catheter in home
 2373 intravenous therapy patients. J Intraven Nurs 1991;14:91-9

2374 259. Harwood IR, Greene LM, Kozakowski-Koch JA and Rasor JS. New peripherally
 2375 inserted midline catheter: a better alternative for intravenous antibiotic therapy in patients
 2376 with cystic fibrosis. Pediatr Pulmonol 1992;12:233-9

2377 260. Mermel LA, Parenteau S and Tow SM. The risk of midline catheterization in
 2378 hospitalized patients. A prospective study. Ann Intern Med 1995;123:841-4

2379 261. Uldall PR, Merchant N, Woods F, Yarworski U and Vas S. Changing subclavian
 2380 haemodialysis cannulas to reduce infection. Lancet 1981;1:1373

Formatted: Swedish (Sweden)

2381 262. Eyer S, Brummitt C, Crossley K, Siegel R and Cerra F. Catheter-related sepsis:
 2382 prospective, randomized study of three methods of long-term catheter maintenance. Crit
 2383 Care Med 1990;18:1073-9

2384 263. Cook D, Randolph A, Kernerman P, et al. Central venous catheter replacement
 2385 strategies: a systematic review of the literature. Crit Care Med 1997;25:1417-24

2386 264. Cobb DK, High KP, Sawyer RG, et al. A controlled trial of scheduled replacement
 2387 of central venous and pulmonary-artery catheters. N Engl J Med 1992;327:1062-8

2388 265. Beathard GA. Management of bacteremia associated with tunneled-cuffed
 2389 hemodialysis catheters. J Am Soc Nephrol 1999;10:1045-9

2390 266. Duszak R, Jr., Haskal ZJ, Thomas-Hawkins C, et al. Replacement of failing tunneled
 2391 hemodialysis catheters through pre-existing subcutaneous tunnels: a comparison of
 2392 catheter function and infection rates for de novo placements and over-the-wire
 2393 exchanges. J Vasc Interv Radiol 1998;9:321-7

2394 267. Robinson D, Suhocki P and Schwab SJ. Treatment of infected tunneled venous
 2395 access hemodialysis catheters with guidewire exchange. Kidney Int 1998;53:1792-4

2396 268. Saad TF. Bacteremia associated with tunneled, cuffed hemodialysis catheters. Am J
 2397 Kidney Dis 1999;34:1114-24

2398 269. Ainsworth SB, Clerihew L and McGuire W. Percutaneous central venous catheters
 2399 versus peripheral cannulae for delivery of parenteral nutrition in neonates. Cochrane
 2400 Database Syst Rev 2007:CD004219

2401 270. Shah PS, Kalyn A, Satodia P, et al. A randomized, controlled trial of heparin versus
 2402 placebo infusion to prolong the usability of peripherally placed percutaneous central

2403 venous catheters (PCVCs) in neonates: the HIP (Heparin Infusion for PCVC) study.
 2404 Pediatrics 2007;119:e284-91

2405 271. Jaar BG, Hermann JA, Furth SL, Briggs W and Powe NR. Septicemia in diabetic
 2406 hemodialysis patients: comparison of incidence, risk factors, and mortality with
 2407 nondiabetic hemodialysis patients. Am J Kidney Dis 2000;35:282-92.

2408 272. Powe NR, Jaar B, Furth SL, Hermann J and Briggs W. Septicemia in dialysis
 2409 patients: incidence, risk factors, and prognosis. Kidney Int 1999;55:1081-90.

2410 273. Hoen B, Paul-Dauphin A, Hestin D and Kessler M. EPIBACDIAL: a multicenter
 2411 prospective study of risk factors for bacteremia in chronic hemodialysis patients. J Am
 2412 Soc Nephrol 1998;9:869-76

2413 274. Blot F, Chachaty E, Raynard B, Antoun S, Bourgain JL and Nitenberg G.
 2414 Mechanisms and risk factors for infection of pulmonary artery catheters and introducer
 2415 sheaths in cancer patients admitted to an intensive care unit. J Hosp Infect 2001;48:289-
 2416 97

2417 275. Kac G, Durain E, Amrein C, Herisson E, Fiemeyer A and Buu-Hoi A. Colonization
 2418 and infection of pulmonary artery catheter in cardiac surgery patients: epidemiology and
 2419 multivariate analysis of risk factors. Crit Care Med 2001;29:971-5

2420 276. Raad I, Umphrey J, Khan A, Truett LJ and Bodey GP. The duration of placement as
 2421 a predictor of peripheral and pulmonary arterial catheter infections. J Hosp Infect
 2422 1993;23:17-26

2423 277. Chen YY, Yen DH, Yang YG, Liu CY, Wang FD and Chou P. Comparison between
 2424 replacement at 4 days and 7 days of the infection rate for pulmonary artery catheters in an
 2425 intensive care unit. Crit Care Med 2003;31:1353-8

2426 278. Boo NY, Wong NC, Zulkifli SS and Lye MS. Risk factors associated with umbilical
 2427 vascular catheter-associated thrombosis in newborn infants. *J Paediatr Child Health*
 2428 1999;35:460-5

2429 279. Garland JS, Buck RK, Maloney P, et al. Comparison of 10% povidone-iodine and
 2430 0.5% chlorhexidine gluconate for the prevention of peripheral intravenous catheter
 2431 colonization in neonates: a prospective trial. *Pediatr Infect Dis J* 1995;14:510-6

2432 280. Krauss AN, Albert RF and Kannan MM. Contamination of umbilical catheters in the
 2433 newborn infant. *J Pediatr* 1970;77:965-9

2434 281. Landers S, Moise AA, Fraley JK, Smith EO and Baker CJ. Factors associated with
 2435 umbilical catheter-related sepsis in neonates. *Am J Dis Child* 1991;145:675-80

2436 282. Cronin WA, Germanson TP and Donowitz LG. Intravascular catheter colonization
 2437 and related bloodstream infection in critically ill neonates. *Infect Control Hosp Epidemiol*
 2438 1990;11:301-8

2439 283. Miller KL, Coen PE, White WJ, Hurst WJ, Achey BE and Lang CM. Effectiveness
 2440 of skin absorption of tincture of I in blocking radioiodine from the human thyroid gland.
 2441 *Health Phys* 1989;56:911-4

2442 284. Ankola PA, Atakent YS. Effect of adding heparin in very low concentration to the
 2443 infusate to prolong the patency of umbilical artery catheters. *Am J Perinatol*
 2444 1993;10:229-32.

2445 285. David RJ, Merten DF, Anderson JC and Gross S. Prevention of umbilical artery
 2446 catheter clots with heparinized infusates. *Dev Pharmacol Ther* 1981;2:117-26.

2447 286. Horgan MJ, Bartoletti A, Polansky S, Peters JC, Manning TJ and Lamont BM.
 2448 Effect of heparin infusates in umbilical arterial catheters on frequency of thrombotic
 2449 complications. J Pediatr 1987;111:774-8.

2450 287. Fletcher MA, Brown DR, Landers S and Seguin J. Umbilical arterial catheter use:
 2451 report of an audit conducted by the Study Group for Complications of Perinatal Care. Am
 2452 J Perinatol 1994;11:94-9

2453 288. Seguin J, Fletcher MA, Landers S, Brown D and Macpherson T. Umbilical venous
 2454 catheterizations: audit by the Study Group for Complications of Perinatal Care. Am J
 2455 Perinatol 1994;11:67-70

2456 289. Loisel DB, Smith MM, MacDonald MG and Martin GR. Intravenous access in
 2457 newborn infants: impact of extended umbilical venous catheter use on requirement for
 2458 peripheral venous lines. J Perinatol 1996;16:461-6

2459 290. Balagtas RC, Bell CE, Edwards LD and Levin S. Risk of local and systemic
 2460 infections associated with umbilical vein catheterization: a prospective study in 86
 2461 newborn patients. Pediatrics 1971;48:359-67

2462 291. Butler-O'Hara M, Buzzard CJ, Reubens L, McDermott MP, DiGrazio W and
 2463 D'Angio CT. A randomized trial comparing long-term and short-term use of umbilical
 2464 venous catheters in premature infants with birth weights of less than 1251 grams.
 2465 Pediatrics 2006;118:e25-35

2466 292. Martin C, Saux P, Papazian L and Gouin F. Long-term arterial cannulation in ICU
 2467 patients using the radial artery or dorsalis pedis artery. Chest 2001;119:901-6

2468 293. Koh DB, Gowardman JR, Rickard CM, Robertson IK and Brown A. Prospective
2469 study of peripheral arterial catheter infection and comparison with concurrently sited
2470 central venous catheters. Crit Care Med 2008;36:397-402

2471 294. Donowitz LG, Marsik FJ, Hoyt JW and Wenzel RP. Serratia marcescens bacteremia
2472 from contaminated pressure transducers. JAMA 1979;242:1749-51

2473 295. Luskin RL, Weinstein RA, Nathan C, Chamberlin WH and Kabins SA. Extended
2474 use of disposable pressure transducers. A bacteriologic evaluation. JAMA 1986;255:916-
2475 20

2476 296. Maki DG, Hassemer CA. Endemic rate of fluid contamination and related
2477 septicemia in arterial pressure monitoring. Am J Med 1981;70:733-8

2478 297. Mermel LA, Maki DG. Epidemic bloodstream infections from hemodynamic
2479 pressure monitoring: signs of the times. Infect Control Hosp Epidemiol 1989;10:47-53

2480 298. Tenold R, Priano L, Kim K, Rourke B and Marrone T. Infection potential of
2481 nondisposable pressure transducers prepared prior to use. Crit Care Med 1987;15:582-3

2482 299. Thomas F, Burke JP, Parker J, et al. The risk of infection related to radial vs femoral
2483 sites for arterial catheterization. Crit Care Med 1983;11:807-12

2484 300. Leroy O, Billiau V, Beuscart C, et al. Nosocomial infections associated with long-
2485 term radial artery cannulation. Intensive Care Med 1989;15:241-6

2486 301. Fisher MC, Long SS, Roberts EM, Dunn JM and Balsara RK. Pseudomonas
2487 maltophilia bacteremia in children undergoing open heart surgery. JAMA

2488 1981;246:1571-4

Formatted: French (France)

2489 302. Stamm WE, Colella JJ, Anderson RL and Dixon RE. Indwelling arterial catheters as
 2490 a source of nosocomial bacteremia. An outbreak caused by *Flavobacterium* Species. N
 2491 Engl J Med 1975;292:1099-102

2492 303. Weinstein RA, Emori TG, Anderson RL and Stamm WE. Pressure transducers as a
 2493 source of bacteremia after open heart surgery. Report of an outbreak and guidelines for
 2494 prevention. Chest 1976;69:338-44

2495 304. Shinozaki T, Deane RS, Mazuzan JE, Jr., Hamel AJ and Hazelton D. Bacterial
 2496 contamination of arterial lines. A prospective study. JAMA 1983;249:223-5

2497 305. Solomon SL, Alexander H, Eley JW, et al. Nosocomial fungemia in neonates
 2498 associated with intravascular pressure-monitoring devices. *Pediatr Infect Dis* 1986;5:680-
 2499 5

2500 306. Weems JJ, Jr., Chamberland ME, Ward J, Willy M, Padhye AA and Solomon SL.
 2501 *Candida parapsilosis* fungemia associated with parenteral nutrition and contaminated
 2502 blood pressure transducers. *J Clin Microbiol* 1987;25:1029-32

2503 307. Villarino ME, Jarvis WR, O'Hara C, Bresnahan J and Clark N. Epidemic of *Serratia*
 2504 *marcescens* bacteremia in a cardiac intensive care unit. *J Clin Microbiol* 1989;27:2433-6

2505 308. Beck-Sague CM, Jarvis WR, Brook JH, et al. Epidemic bacteremia due to
 2506 *Acinetobacter baumannii* in five intensive care units. *Am J Epidemiol* 1990;132:723-33

2507 309. Rijnders BJ, Van Wijngaerden E, Wilmer A and Peetermans WE. Use of full sterile
 2508 barrier precautions during insertion of arterial catheters: a randomized trial. *Clin Infect*
 2509 *Dis* 2003;36:743-8

2510 310. Scheer B, Perel A and Pfeiffer UJ. Clinical review: complications and risk factors of
 2511 peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and
 2512 intensive care medicine. *Crit Care* 2002;6:199-204

2513 311. Lorente L, Santacreu R, Martin MM, Jimenez A and Mora ML. Arterial catheter-
 2514 related infection of 2,949 catheters. *Crit Care* 2006;10:R83

2515 312. Furfaro S, Gauthier M, Lacroix J, Nadeau D, Lafleur L and Mathews S. Arterial
 2516 catheter-related infections in children. A 1-year cohort analysis. *Am J Dis Child*
 2517 1991;145:1037-43

2518 313. Gillies D, O'Riordan L, Wallen M, Morrison A, Rankin K and Nagy S. Optimal
 2519 timing for intravenous administration set replacement. *Cochrane Database Syst Rev*
 2520 2005:CD003588

2521 314. Sitges-Serra A, Linares J, Perez JL, Jaurrieta E and Lorente L. A randomized trial on
 2522 the effect of tubing changes on hub contamination and catheter sepsis during parenteral
 2523 nutrition. *JPEN J Parenter Enteral Nutr* 1985;9:322-5

2524 315. Snyderman DR, Donnelly-Reidy M, Perry LK and Martin WJ. Intravenous tubing
 2525 containing burettes can be safely changed at 72 hour intervals. *Infect Control* 1987;8:113-
 2526 6

2527 316. Josephson A, Gombert ME, Sierra MF, Karanfil LV and Tansino GF. The
 2528 relationship between intravenous fluid contamination and the frequency of tubing
 2529 replacement. *Infect Control* 1985;6:367-70

2530 317. Melly MA, Meng HC and Schaffner W. Microbiol growth in lipid emulsions used in
 2531 parenteral nutrition. *Arch Surg* 1975;110:1479-81

2532 318. Mershon J, Nogami W, Williams JM, Yoder C, Eitzen HE and Lemons JA.
 2533 Bacterial/fungal growth in a combined parenteral nutrition solution. JPEN J Parenter
 2534 Enteral Nutr 1986;10:498-502

2535 319. Gilbert M, Gallagher SC, Eads M and Elmore MF. Microbial growth patterns in a
 2536 total parenteral nutrition formulation containing lipid emulsion. JPEN J Parenter Enteral
 2537 Nutr 1986;10:494-7

2538 320. Maki DG, Martin WT. Nationwide epidemic of septicemia caused by contaminated
 2539 infusion products. IV. Growth of microbial pathogens in fluids for intravenous infusions.
 2540 J Infect Dis 1975;131:267-72

2541 321. Bennett SN, McNeil MM, Bland LA, et al. Postoperative infections traced to
 2542 contamination of an intravenous anesthetic, propofol. N Engl J Med 1995;333:147-54

2543 322. Rickard CM, Lipman J, Courtney M, Siversen R and Daley P. Routine changing of
 2544 intravenous administration sets does not reduce colonization or infection in central
 2545 venous catheters. Infect Control Hosp Epidemiol 2004;25:650-5

2546 323. Hanna HA, Raad I. Blood products: a significant risk factor for long-term catheter-
 2547 related bloodstream infections in cancer patients. Infect Control Hosp Epidemiol
 2548 2001;22:165-6.

2549 324. Saiman L, Ludington E, Dawson JD, et al. Risk factors for Candida species
 2550 colonization of neonatal intensive care unit patients. Pediatr Infect Dis J 2001;20:1119-
 2551 24.

2552 325. Avila-Figueroa C, Goldmann DA, Richardson DK, Gray JE, Ferrari A and Freeman
 2553 J. Intravenous lipid emulsions are the major determinant of coagulase-negative

Formatted: Swedish (Sweden)

Formatted: Swedish (Sweden)

2554 staphylococcal bacteremia in very low birth weight newborns. *Pediatr Infect Dis J*
 2555 1998;17:10-7.

2556 326. Crocker KS, Noga R, Filibeck DJ, Krey SH, Markovic M and Steffee WP. Microbial
 2557 growth comparisons of five commercial parenteral lipid emulsions. *J Parenter Enteral*
 2558 *Nutr* 1984;8:391-5

2559 327. Jarvis WR, Highsmith AK. Bacterial growth and endotoxin production in lipid
 2560 emulsion. *J Clin Microbiol* 1984;19:17-20

2561 328. Arduino MJ, Bland LA, Danzig LE, McAllister SK and Aguero SM. Microbiologic
 2562 evaluation of needleless and needle-access devices. *Am J Infect Control* 1997;25:377-80

2563 329. Brown JD, Moss HA and Elliott TS. The potential for catheter microbial
 2564 contamination from a needleless connector. *J Hosp Infect* 1997;36:181-9

2565 330. Cookson ST, Ihrig M, O'Mara EM, et al. Increased bloodstream infection rates in
 2566 surgical patients associated with variation from recommended use and care following
 2567 implementation of a needleless device. *Infect Control Hosp Epidemiol* 1998;19:23-7

2568 331. Seymour VM, Dhallu TS, Moss HA, Tebbs SE and Elliot TS. A prospective clinical
 2569 study to investigate the microbial contamination of a needleless connector. *J Hosp Infect*
 2570 2000;45:165-8

2571 332. Luebke MA, Arduino MJ, Duda DL, et al. Comparison of the microbial barrier
 2572 properties of a needleless and a conventional needle-based intravenous access system.
 2573 *Am J Infect Control* 1998;26:437-41

2574 333. McDonald LC, Banerjee SN and Jarvis WR. Line-associated bloodstream infections
 2575 in pediatric intensive-care-unit patients associated with a needleless device and
 2576 intermittent intravenous therapy. *Infect Control Hosp Epidemiol* 1998;19:772-7

Formatted: French (France)

2577 334. Mendelson MH, Short LJ, Schechter CB, et al. Study of a needleless intermittent
 2578 intravenous-access system for peripheral infusions: analysis of staff, patient, and
 2579 institutional outcomes. Infect Control Hosp Epidemiol 1998;19:401-6

2580 335. Do AN, Ray BJ, Banerjee SN, et al. Bloodstream infection associated with
 2581 needleless device use and the importance of infection-control practices in the home health
 2582 care setting. J Infect Dis 1999;179:442-8

2583 336. Rupp ME, Sholtz LA, Jourdan DR, et al. Outbreak of bloodstream infection
 2584 temporally associated with the use of an intravascular needleless valve. Clin Infect Dis
 2585 2007;44:1408-14

2586 337. Salgado CD, Chinnes L, Paczesny TH and Cantey JR. Increased rate of catheter-
 2587 related bloodstream infection associated with use of a needleless mechanical valve device
 2588 at a long-term acute care hospital. Infect Control Hosp Epidemiol 2007;28:684-8

2589 338. Maragakis LL, Bradley KL, Song X, et al. Increased catheter-related bloodstream
 2590 infection rates after the introduction of a new mechanical valve intravenous access port.
 2591 Infect Control Hosp Epidemiol 2006;27:67-70

2592 339. Field K, McFarlane C, Cheng AC, et al. Incidence of catheter-related bloodstream
 2593 infection among patients with a needleless, mechanical valve-based intravenous
 2594 connector in an Australian hematology-oncology unit. Infect Control Hosp Epidemiol
 2595 2007;28:610-3

2596 340. Inoue Y, Nezu R, Matsuda H, et al. Prevention of catheter-related sepsis during
 2597 parenteral nutrition: effect of a new connection device. J Parenter Enteral Nutr
 2598 1992;16:581-5

Formatted: French (France)

Formatted: Swedish (Sweden)

2599 341. Yebenes JC, Vidaur L, Serra-Prat M, et al. Prevention of catheter-related
 2600 bloodstream infection in critically ill patients using a disinfectable, needle-free connector:
 2601 a randomized controlled trial. *Am J Infect Control* 2004;32:291-5

2602 342. Casey AL, Burnell S, Whinn H, Worthington T, Faroqui MH and Elliott TS. A
 2603 prospective clinical trial to evaluate the microbial barrier of a needleless connector. *J*
 2604 *Hosp Infect* 2007;65:212-8

2605 343. Casey AL, Worthington T, Lambert PA, Quinn D, Faroqui MH and Elliott TS. A
 2606 randomized, prospective clinical trial to assess the potential infection risk associated with
 2607 the PosiFlow needleless connector. *J Hosp Infect* 2003;54:288-93

2608 344. Esteve F, Pujol M, Limon E, et al. Bloodstream infection related to catheter
 2609 connections: a prospective trial of two connection systems. *J Hosp Infect* 2007;67:30-4

2610 345. Yebenes JC, Delgado M, Sauca G, et al. Efficacy of three different valve systems of
 2611 needle-free closed connectors in avoiding access of microorganisms to endovascular
 2612 catheters after incorrect handling. *Crit Care Med* 2008;36:2558-61

2613 346. Menyhay SZ, Maki DG. Disinfection of needleless catheter connectors and access
 2614 ports with alcohol may not prevent microbial entry: the promise of a novel antiseptic-
 2615 barrier cap. *Infect Control Hosp Epidemiol* 2006;27:23-7

2616 347. Herruzo-Cabrera R, Garcia-Caballero J, Vera-Cortes ML, et al. Growth of
 2617 microorganisms in parenteral nutrient solutions. *Am J Hosp Pharm* 1984;41:1178-80

2618 348. ASHP guidelines on quality assurance for pharmacy-prepared sterile products.
 2619 American Society of Health System Pharmacists. *Am J Health Syst Pharm* 2000;57:1150-
 2620 69

Formatted: Italian (Italy)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

2621 349. Green KA, Mustachi B, Schoer K, Moro D, Blend R and McGeer A. Gadolinium-
2622 based MR contrast media: potential for growth of microbial contaminants when single
2623 vials are used for multiple patients. *AJR Am J Roentgenol* 1995;165:669-71

Formatted: German (Germany)

2624 350. Salzman MB, Isenberg HD and Rubin LG. Use of disinfectants to reduce microbial
2625 contamination of hubs of vascular catheters. *J Clin Microbiol* 1993;31:475-9

Formatted: Italian (Italy)

2626 351. Arrington ME, Gabbert KC, Mazgaj PW and Wolf MT. Multidose vial
2627 contamination in anesthesia. *Aana J* 1990;58:462-6

2628 352. Plott RT, Wagner RF, Jr. and Tying SK. Iatrogenic contamination of multidose
2629 vials in simulated use. A reassessment of current patient injection technique. *Arch*
2630 *Dermatol* 1990;126:1441-4

2631 353. Didier ME, Fischer S and Maki DG. Total nutrient admixtures appear safer than
2632 lipid emulsion alone as regards microbial contamination: growth properties of microbial
2633 pathogens at room temperature. *JPEN J Parenter Enteral Nutr* 1998;22:291-6

Formatted: Swedish (Sweden)

2634 354. Barrett BB, Andersen JW and Anderson KC. Strategies for the avoidance of
2635 bacterial contamination of blood components. *Transfusion* 1993;33:228-33

2636 355. Blajchman MA. Reducing the risk of bacterial contamination of cellular blood
2637 components. *Dev Biol (Basel)* 2000;102:183-93

2638 356. Roth VR, Arduino MJ, Nobiletti J, et al. Transfusion-related sepsis due to *Serratia*
2639 *liquefaciens* in the United States. *Transfusion* 2000;40:931-5

2640 357. Wagner SJ, Friedman LI and Dodd RY. Transfusion-associated bacterial sepsis. *Clin*
2641 *Microbiol Rev* 1994;7:290-302

2642 358. Longfield RN, Smith LP, Longfield JN, Coberly J and Cruess D. Multiple-dose
 2643 vials: persistence of bacterial contaminants and infection control implications. *Infect*
 2644 *Control* 1985;6:194-9

2645 359. Henry B, Plante-Jenkins C and Ostrowska K. An outbreak of *Serratia marcescens*
 2646 associated with the anesthetic agent propofol. *Am J Infect Control* 2001;29:312-5.

2647 360. Grohskopf LA, Roth VR, Feikin DR, et al. *Serratia liquefaciens* bloodstream
 2648 infections from contamination of epoetin alfa at a hemodialysis center. *N Engl J Med*
 2649 2001;344:1491-7.

2650 361. Thompson ND, Perz JF, Moorman AC and Holmberg SD. Nonhospital healthcare-
 2651 associated hepatitis B and C virus transmission: United States, 1998-2008. *Ann Intern*
 2652 *Med* 2009;150:33-9

2653 362. Costello JM, Morrow DF, Graham DA, Potter-Bynoe G, Sandora TJ and Laussen
 2654 PC. Systematic intervention to reduce central line-associated bloodstream infection rates
 2655 in a pediatric cardiac intensive care unit. *Pediatrics* 2008;121:915-23

2656 363. Frankel HL, Crede WB, Topal JE, Roumanis SA, Devlin MW and Foley AB. Use of
 2657 corporate Six Sigma performance-improvement strategies to reduce incidence of
 2658 catheter-related bloodstream infections in a surgical ICU. *J Am Coll Surg* 2005;201:349-
 2659 58

2660 364. Galpern D, Guerrero A, Tu A, Fahoum B and Wise L. Effectiveness of a central line
 2661 bundle campaign on line-associated infections in the intensive care unit. *Surgery*
 2662 2008;144:492-5; discussion 495

2663 365. McKee C, Berkowitz I, Cosgrove SE, et al. Reduction of catheter-associated
 2664 bloodstream infections in pediatric patients: experimentation and reality. *Pediatr Crit*
 2665 *Care Med* 2008;9:40-6

2666 366. Pronovost PJ, Berenholtz SM and Goeschel CA. Improving the quality of
 2667 measurement and evaluation in quality improvement efforts. *Am J Med Qual*
 2668 2008;23:143-6

2669 367. Safdar N, Maki DG. Lost in translation. *Infect Control Hosp Epidemiol* 2006;27:3-7

2670 368. Warren DK, Yokoe DS, Climo MW, et al. Preventing catheter-associated
 2671 bloodstream infections: a survey of policies for insertion and care of central venous
 2672 catheters from hospitals in the prevention epicenter program. *Infect Control Hosp*
 2673 *Epidemiol* 2006;27:8-13

2674 369. Krein SL, Hofer TP, Kowalski CP, et al. Use of central venous catheter-related
 2675 bloodstream infection prevention practices by US hospitals. *Mayo Clin Proc*
 2676 2007;82:672-8

2677 370. Lobo RD, Levin AS, Gomes LM, et al. Impact of an educational program and policy
 2678 changes on decreasing catheter-associated bloodstream infections in a medical intensive
 2679 care unit in Brazil. *Am J Infect Control* 2005;33:83-7

2680 371. Marschall J, Leone C, Jones M, Nihill D, Fraser VJ and Warren DK. Catheter-
 2681 associated bloodstream infections in general medical patients outside the intensive care
 2682 unit: a surveillance study. *Infect Control Hosp Epidemiol* 2007;28:905-9

2683 372. Rosenthal VD, McCormick RD, Guzman S, Villamayor C and Orellano PW. Effect
 2684 of education and performance feedback on handwashing: the benefit of administrative
 2685 support in Argentinean hospitals. *Am J Infect Control* 2003;31:85-92

Formatted: German (Germany)

2686 373. Gastmeier P, Geffers C. Prevention of catheter-related bloodstream infections:
2687 analysis of studies published between 2002 and 2005. J Hosp Infect 2006;64:326-35
2688 374. Shapey IM, Foster MA, Whitehouse T, Jumaa P and Bion JF. Central venous
2689 catheter-related bloodstream infections: improving post-insertion catheter care. J Hosp
2690 Infect 2009;71:117-22
2691
2692

DRAFT